

DIET HISTORY AND BIRTH WEIGHT RELATIONSHIP AMONG SAUDI PREGNANT WOMEN

Ahmed A. Al-Shoshan¹

ABSTRACT

Objective: The aim of this study was to see the relationship between food intake pattern and its effect on birth weight.

Subjects and Methods: 1771 Saudi pregnant women who were admitted for delivery at the government maternity hospitals in Riyadh city of Saudi Arabia who gave birth to singleton live-born babies were interviewed immediately after delivery.

Results: During the pregnancy the average energy and protein intakes were 1815.7±624Cal. and 86±30.9g, respectively. Intakes of vitamins A and B12 supply 90% and 92% of RDA for each nutrient, respectively. The intake of 94%, 91% and 90% of women were below 85% of RDA for folate, iron and vitamin B1, respectively.

Conclusions: Mothers of heavier infants have significantly higher average intake of most nutrients. Intake level of individual nutrients correlates significantly with NAR index which parallel the findings for individual nutrient intakes and was the only variable in the regression equation calculated.

KEY WORDS: Pregnancy, Maternal nutrition, Birth weight.

Pak J Med Sci April 2007 Vol. 23 No. 2 176-181

INTRODUCTION

Pregnancy is a natural phenomenon during which women encounter a wide range of internal physical and emotional changes. Quantification of the impact of nutrition on human reproductive performance is important to planners and to justify investment in maternal nutrition programs. This impact should reside in the current pregnancy for intervention to be effective.¹ Pregnancy and the outcome growth are affected by maternal and various environmental and demographic factors.² Growth in the intrauterine phase is

dependent upon genetic factors and environmental support. Placental factors include vascular abnormalities, placental hormones and hypoxia. Chromosomal abnormalities or syndromes within the fetus itself influence fetal growth.³ Because birth weight is often used as an indicator of the successful outcome of pregnancy, studies have attempted to explain variability in birth weight attributable to these factors.⁴⁻⁸ The positive relationship between maternal nutrition and birth weight has been reported under acute starvation⁹⁻¹¹ but less clear with moderate levels of malnutrition.¹²⁻¹⁴ In general, some studies have included dietary factors^{4,15} and/or examine overall dietary quality as an independent variable.¹⁶ Assessment of dietary intake during pregnancy is important because it is well established that both nutrient deficiencies and excesses can have adverse effects on pregnancy outcome.¹⁷

Among groups of pregnant women in Saudi Arabia, deficiency of Vitamin D, nutrition knowledge and food habits has been studied.¹⁸⁻²¹ The effect on birth weight in

1. Ahmed A. Al-Shoshan
Associate professor
Food Science and Nutrition Department
College of Food and Agricultural Sciences
King Saud University,
Riyadh - Saudi Arabia.

Correspondence

Ahmed A. Al-Shoshan
E-mail: Draahmedan@hotmail.com

* Received for Publication: December 28, 2006

* Revision Received: January 5, 2007

* Final Revision Accepted: March 11, 2007

relation to other factors was also investigated.²² However, literature on the nutritional status of Saudi pregnant women is scarce, and the relationship between maternal nutrition and the birth weight of infants demand further attention. This study was designed to measure the dietary intake and nutrition adequacy of Saudi women throughout pregnancy and to relate the birth weight of infant born to these nutritional factors.

SUBJECTS AND METHODS

A total of 1771 Saudi pregnant women were interviewed soon after delivery for their food intake during the current pregnancy. The data were collected 24 hours after delivery from women delivered at the government maternity hospitals in Riyadh city. A structured food frequency recall questionnaire was used in which most common food items were included. Additional items of food were also incorporated according to the mother's consumption. The foods were grouped into exchange lists, their nutrient content analyzed and food scores calculated with a maximum score of 25 for each of the food groups. All figures were compared to the Recommended Dietary Allowances (RDAs) for pregnant women.²³ The data was then used to estimate intake levels of energy, twelve nutrients and five food groups, and to calculate a nutrient adequacy ratio index (NAR). Nutrients estimated were protein, vitamin A, vitamin C, vitamin B1, vitamin B2, vitamin B6, vitamin B12, calcium, phosphorus, Iron and folate. The food groups were milk, fruit, vegetable, cereal and meat group. The NAR index for each pregnant women was obtained by converting intake to percentage of RDA for 12 nutrients, truncating amounts over 100% so that excessive amounts of some nutrients would not affect the index, and then averaging the percentages, a procedure similar to that used by Philipps and Johnson.¹⁵ Data on neonatal outcome was obtained from the babies' medical records for singleton live-born babies.

To relate infant birth weight to food intake and nutrition adequacy of expecting mother,

pregnant women under study were grouped according to their intake above and below the RDA level for each nutrient. The resulting average birth weight was then calculated. Average intake from each nutrient and food group was also determined for the mothers having babies with weight below and above 2500g.

Data coding was carried out using commercial Dbase IV and the subsequent analyses were performed by the use of SPSS/PC+ statistical package.

RESULTS

The average of daily intakes, intake as a percentage of the RDA and the percentage of women with intake below 85% of the RDA for energy and selected nutrients are presented in Table-I. Average energy and protein intake were 1815.7±624 Cal and 86±30.9g which provided around 65.7% and 96.0% of the recommended levels of each respectively. Pregnant women had an intake levels that supply 90.5% and 92% of RDA for vitamins A and B12 respectively. Folate, Iron and vitamin B1 intakes were below 85% of RDA for 94%, 91% and 89.6% of women respectively. The average intake was less than the RDA for zinc followed by magnesium, and meet the recommended for niacin. The average NAR index was 69.17% of total RDA for the 12 nutrients included. Average score of intake of food groups and the percentage of women with intake below 85% of the RDA exchange for each group are shown in Table-II. The consumption of milk, meat, and vegetable groups fall below 85% of the RDA for 85.7%, 79.4%, and 53.3% of the pregnant women respectively, and meet the recommended exchanges among 81.8% and 84.2% women for fruit and cereal groups respectively. The NAR index parallel the findings for individual nutrient intakes.

Table-III presents the average of infant birth weight for women with intake below and above RDA levels for each nutrient. Average birth weight was slightly higher among mothers with intake above 85% of the recommended, compared to those with less intake. The birth

Table-I: Average daily intakes; the average intake as a percentage of the RDA and the percentage of women with intake below 85% of the RDA for energy and selected nutrients.

Nutrient	Mean \pm SD	RDA	intakeof RDA ^a	% below 85% RDA ^b
Calorie (Cal)	1815.74 \pm 624.06	2700	65.70 \pm 19.58	54.5%
Protein (g)	85.99 \pm 30.90	60	96.03 \pm 11.04	9.8%
Vitamin A (Mg RE)	1034.96 \pm 480.02	800	90.45 \pm 18.19	29.3%
Vitamin C (mg)	75.87 \pm 43.07	70	82.40 \pm 22.92	40.5%
Folate (Mg)	77.75 \pm 26.74	400	19.43 \pm 6.69	94.0%
Vitamin B1 (mg)	71.00 \pm 0.27	1.5	46.84 \pm 17.48	89.6%
Vitamin B2 (mg)	1.77 \pm 0.80	1.6	86.79 \pm 19.33	29.5%
Vitamin B6 (mg)	1.27 \pm 0.51	2.2	56.85 \pm 21.41	72.4%
Vitamin B12 (Mg)	3.12 \pm 1.58	2.2	91.97 \pm 16.83	19.3%
Calcium (mg)	897.17 \pm 450.68	1200	68.46 \pm 25.10	72.2%
Phosphorus (mg)	1253.32 \pm 467.96	1200	87.63 \pm 17.73	37.9%
Iron (mg)	11.22 \pm 4.19	30	37.37 \pm 13.97	91.1%
Magnesium (mg)	246.33 \pm 97.74	320		
Zinc (mg)	6.49 \pm 2.79	15		
Niacin (mg NE)	17.81 \pm 7.85	17		
Carbohydrate (g)	250.92 \pm 101.04			
Total Fat (g)	43.49 \pm 18.37			
Cholesterol (mg)	286.01 \pm 151.92			
NAR index	69.17 \pm 13.70			

^a The average of intake among pregnant as a percentage of RDA for each nutrient.

^b The percentage with intakes below 85% of the RDA.

weight difference was significant with higher intake of vitamin A, vitamin C, phosphorus ($P \leq 0.001$), vitamin B2 ($P \leq 0.05$), and was insignificant in respect to energy and other nutrients including protein intake. Means of intake for mothers with birth weight below and above 2500g for nutrient and food groups are presented in Table-IV and Table-V respectively. Mothers with infant weight over 2500g show a significant higher intake of energy and nutrients, except Niacin. Their RNA index was $69.50 \pm 13.47\%$ which is significantly higher than that for those with smaller infant weight, 65.34 ± 15.72 .

DISCUSSION

In a simple correlation test, intake level of individual nutrients correlates significantly with overall adequacy measured by RNA index ($P \leq 0.001$). Both show no correlation with the resulting birth weight of infants. In multiple regression stepwise method, to predict birth weight, the total R^2 was 0.069, with RNA index being the only variable in the equation.

An appropriate eating pattern is essential throughout the childbearing years and during

pregnancy to ensure a healthy pregnancy and baby.²⁴ In a country such as Saudi Arabia where food is easily available, nutritional status of women during the course of pregnancy is expected to be compatible to those in other societies with similar standards of living. In this study the averages of intake of Saudi pregnant women for energy and selected nutrients were compared to the recommended daily allowance.

Pregnant women in this study found to consume on average $65.7 \pm 20\%$ of the recommended calories which left more than 54.5% of them below two third of RDA level. Almost all pregnant women included had a level of intake far below the recommended from

Table-II: Scores of intake of food groups and the percentage of women with intake < 85 % of the RDA exchanges.

Food groups	Mean+SD ^a	RDA Exchange	% women below 85 % of RDA
Milk	15.92 \pm 6.95	4-5	85.7
Fruit	23.26 \pm 4.23	2-3	18.2
Vegetable	21.11 \pm 2.51	2-3	53.3
Cereal	23.62 \pm 3.67	4-6	15.8
Meat	19.24 \pm 6.15	7-8	79.4

^a Maximum score is 25.

Table-III: means of birth weight for women with intake < and \geq RDA levels for each nutrient.

Nutrient	Birth Weight Mean \pm SD		P Value
	Group 1 ^a	Group 2 ^b	
Calorie	3194.83 \pm 595.21	3227.14 \pm 521.22	0.22
Protein	3181.42 \pm 615.30	3212.56 \pm 557.00	0.48
Vitamin A	3151.23 \pm 609.62	3233.71 \pm 540.67	0.001
Vitamin C	3163.82 \pm 599.59	3240.67 \pm 534.42	0.001
Folate	3209.53 \pm 562.84		
Vitamin B1	3207.63 \pm 566.61	3250.63 \pm 474.98	0.50
Vitamin B2	3164.90 \pm 599.98	3228.24 \pm 545.68	0.02
Vitamin B6	3204.00 \pm 574.57	3224.04 \pm 531.11	0.50
Vitamin B12	3177.60 \pm 581.79	3217.16 \pm 558.15	0.24
Calcium	3209.38 \pm 580.06	3209.91 \pm 516.04	
Phosphorus	3162.86 \pm 598.21	3238.01 \pm 538.41	0.003
Iron	3209.19 \pm 564.54	3246.87 \pm 330.38	0.78

^a Birth weight for group of mothers with intake < 85% of the RDA.

^b Birth weight for group of mothers with intake \geq 85% of the RDA.

folate, iron and vitamin B1. Meat, milk and vegetable groups were below the consumption level expected.

In interpreting the insufficient intake observed, certain factors are worth mentioning. First, this study was carried out at government hospitals in which a larger

percentage of deliveries are that for women of low and middle socioeconomic levels. Second, the RNA index value obtained parallel the findings for individual nutrient intakes which may support its use as a better overall measure to test the adequacy of mother's diet. In this regard, the quality of the mother's diet expressed as NAR index determined by this study is somewhat comparable to that of American women, 74.7 \pm 12.8.¹⁷

Birth weight was slightly affected by level of intake of most nutrients and food groups (Tables III-V). Higher intake and higher RNA index, on average, resulted on heavier babies and smaller babies were born to mothers who had less intake of nutrients and food groups during pregnancy. However, Hutcheon, et al²⁵ related higher birth weights to increased maternal age, pre-pregnancy body mass index, weight gain during pregnancy and carbohydrate intake. Such relative adequacy of nutrients and food choice of food groups as found in this study did not, on average, determine (insignificant figures) the positive effect on birth weight. The absence of a meaningful positive correlation between either individual nutrients

Table IV: Means of intake for mothers with birth weight < and \geq 2500g for selected nutrients.

Nutrient	Intake (Mean \pm SD)		P Value
	Group 1 ^a	Group 2 ^b	
Calorie (Cal)	1692.28 \pm 593.34	1826.51 \pm 625.69	0.01
Protein (g)	79.69 \pm 28.94	86.54 \pm 31.01	0.01
Vitamin A (Mg RE)	895.93 \pm 441.17	1047.09 \pm 481.48	0.004
Vitamin C (mg)	69.36 \pm 40.17	76.43 \pm 43.27	0.05
Folate (Mg)	72.97 \pm 27.66	78.16 \pm 26.62	0.02
Vitamin B1 (mg)	0.64 \pm 00.24	0.71 \pm 0.27	0.00
Vitamin B2 (mg)	1.57 \pm 00.69	1.79 \pm 0.80	0.001
Vitamin B6 (mg)	1.17 \pm 00.50	1.27 \pm 0.51	0.02
Vitamin B12 (Mg)	2.80 \pm 01.27	3.15 \pm 1.60	0.01
Calcium (mg)	809.70 \pm 413.75	905.89 \pm 453.05	0.01
Phosphorus (mg)	1150.28 \pm 443.13	1262.31 \pm 469.11	0.003
Iron (mg)	10.27 \pm 3.99	11.29 \pm 4.20	0.001
Magnesium (mg)	225.12 \pm 94.06	248.17 \pm 97.86	0.001
Zinc (mg)	5.94 \pm 02.45	6.53 \pm 2.80	0.01
Niacin (mg NE)	16.95 \pm 07.29	17.88 \pm 7.89	0.17
Carbohydrate (g)	234.37 \pm 93.10	252.36 \pm 101.60	0.04
Total Fat (g)	39.68 \pm 17.04	43.81 \pm 18.45	0.003
Cholesterol (mg)	259.84 \pm 143.54	288.29 \pm 152.46	0.03
NAR index	65.34 \pm 15.72	69.50 \pm 13.47	0.002

^a Mothers with newborn birth weight < 2500g.

^b Mothers with newborn birth weight \geq 2500g.

Table-V: means of intake for mothers with birth weight < and \geq 2500 g for food groups.

Food groups	Mean \pm SD	Intake score		P Value
		Mean \pm SD		
		Group 1 ^a	Group 2 ^b	
Milk	15.92 \pm 6.95	14.80 \pm 7.27	16.02 \pm 6.92	0.04
Fruit				
& Veg	23.26 \pm 4.23	22.56 \pm 4.85	23.32 \pm 4.17	0.04
Cereal	23.62 \pm 3.67	23.26 \pm 3.94	23.65 \pm 3.65	0.21
Meat	19.24 \pm 6.15	18.14 \pm 6.48	19.33 \pm 6.11	0.02

^a Mothers with newborn birth weight <2500g.

^b Mothers with newborn birth weight \geq 2500g.

or food group intakes and birth weight indicate that birth weight was not determinantal factor among mothers, regardless of their level of intake. The NAR index being the only variable in the regression equation calculated, support its use as a better determinant of birth weight. Studies investigating the association between women diet and the outcome of pregnancy, reported inconsistent results.²⁶ In their study, Seiler and Fox²⁷ found no significant correlation between dietary score and birth weight. This absence of clear relationship between birth weight and food intake during pregnancy on those studies could be due in part to the fact that most of the women of the studied populations were relatively well nourished. In this study, the average intake was insufficient indicator of birth weight for both groups. Furthermore, the number of women included in this study, having an outcome with birth weight <2500g was 140 compared to 1631 women who gave birth weight \geq 2500g. Although there is evidence for the importance of maternal nutrition as a determinant of the course and outcome of pregnancy, inconsistent results being reported from attempts to find direct correlation between diet and birth weight of infants.^{1,17} There appears to be a minimal level of nutrients which must be available in order to obtain adequate birth weight. However, above this minimum level, pregnant women can adapt themselves to a wide variety of food intake, both in quantity and quality, without affecting birth weight.²⁸

In conclusion, the observed low intake of energy may divert the use of sufficient ingested

protein as a source of energy and in turn may make such diet inadequate even for protein. The amounts of iron and folate provided in the pregnant women's diet, could not meet the greatly increased requirements during this state of life. And even though those nutrients are widely supplemented for pregnant women around the country, the observed magnitude of their deficiency should raise a health concern and worth deeper investigation. In this respect, Turner, et al²⁹ concluded that the participants in their study did not consume adequate amounts of iron from food to meet the needs of pregnancy, and therefore iron supplementation is warranted in that population. The average diet (as revealed in this study) is deficient in important nutrients and the importance of micronutrients, specially of plant source, is to be addressed along with vitamin D and calcium which has been proven by others as deficient.^{23,18}

The upraise for nutrition awareness programs among pregnant women is recommended. Food preferences should be encouraged to accommodate suitable sources in term of adequate energy and micronutrient intakes. The fact that the average of low birth weight rate is within an acceptable figure should not lead to overlooking of other less observed signs and effects of nutrient deficiencies. However, there could be other factors responsible for low birth weight in spite of maternal nutrition.

ACKNOWLEDGEMENT

The author would like to acknowledge the unfailing support and assistance provided by all concerned personal and employees of the Health Ministry and The Nutrition Department in Riyadh.

REFERENCES

1. Bergner L, Susser MW. Low birth weight and prenatal nutrition: an interpretative review. *Pediatrics* 1970;46:946-56.
2. MUSAIGER AO. Factors associated with birthweight in Bahrain. *J Trop Medicine Hygiene* 1985;88:31-6.
3. GOKHALE R, KIRSCHNER BS. Assessment of growth and Nutrition. *Best practice and Research Clinical Gastroenterology* 2003;17(2):153-62.

4. Higgins AC. Montreal diet dispensary study Nutritional Supplementation and the outcome of pregnancy. Proceedings of a Workshop. Washington DC, Natl Acad Sci-Natl Res Council 2003; pp 93-111.
5. Rush D. Examination of the relationship between birth weight, cigarette smoking during pregnancy and maternal weight gain. *J Obstet Gynaecol Br Commonw* 1974;81:746-51.
6. O'sullivan JB, Gellis SS, Tenney BO. Aspects of birth weight and its influencing variables. *Am J Obstet Gynecol* 1965;92:1023-7.
7. Weiss W, Jackson EC. Maternal factors affecting birth weight. In: *Prenatal Factors Affecting Human Development*. PAHO Sci, Washington DC, Pan American Health Organization 1969;(185) pp 54-55.
8. Naylor AF, Myrlandhopouloss NC. The relation of ethnic and selected socio-economic factors to human birth weight. *Ann Hum Genet* 1967;31:71-7.
9. Antonov AN. Children born during the siege of Leningrad in 1942. *J Pediatr* 1947;30:250-5.
10. South CA. Effects of maternal undernutrition upon the newborn infant in Holland (1944-45). *J Pediatr* 1947;30:229-34.
11. Gruenwald P, Funakawa IL. Influence of environmental factors on fetal growth in man. *Lancet* 1967;1:1026-30.
12. Chase HC. Infant mortality and weight at birth: 1960-United States cohort. *Am J Public Health* 1969;59:1618-25.
13. Mate LJ, Urrutia JJ. Influence of recurrent infections on nutrition and growth of children in Guatemala. *Am J Clin Nutr* 1972;25:1267-72.
14. McCance RA, Widdowson EM. Nutrition and growth. *Proc R Soc Biol* 1975; 156:326-34.
15. Philipps C, Johnson NE. The impact of quality of diet and other factors on birth weight of infants. *Am J Clin Nutr* 1977;30:215-25.
16. Lechtig A, Habicht JP, Delgado H. Effect of food supplementation during pregnancy on birth weight. *J Pediatr* 1975;56:508-19.
17. Worthington RB. The role of maternal nutrition in the prevention of birth defects. *J Amer Dietetic Assoc* 1975;97:184-5.
18. Serenius F, Elidrissy AT. Vitamin D nutrition in pregnant women at term and in newly born babies in Saudi Arabia. *J Clin Pathol* 1984;37:444-7.
19. Mansour AA, Hassan SA. Factors that influence women's nutrition knowledge in Saudi Arabia. *Health Care Women Int* 1994;15:213-23.
20. Al-Kanhal MA, Bani IA. Food habits during pregnancy among Saudi women. *Int. J Vitam Nutr Res* 1995;65:206-10.
21. Lawoyin TO, Ani F. Epidemiological aspects of pre-eclampsia in Saudi Arabia. *East Afr Med J* 1996;73:404-6.
22. Rasheed P, Rahman J. Predictors of Saudi Birth weights: A Multiple Regression Analysis. *Saudi Med J* 1995;16:23-9.
23. *Recommended Dietary Allowances, 8th ed.* Washington DC, Natl Acad. Sci.-Natl Res Council Publ 1974; pp 2216.
24. Pick ME, Edwards M, Moreau D, Ryan EA. Assessment of diet quality in Pregnant women using the healthy eating index. *J Amer Dietetic Assoc* 2005;105:240-6.
25. Hutcheon JA, Platt RW, Meltzer SJ, Egeland GM. Is birth weight modified during pregnancy? Using sibling differences to understand the impact of blood glucose, obesity, and maternal weight gain in gestational diabetes. *Amer J Obstet Gynecol* 2005;195:488-94.
26. Wahid MA, Fathi SA. Nutrition and the unborn baby. *Ric Clin Lab* 1987;17:199-206.
27. Seiler JA, Fox HM. Adolescent pregnancy: association of dietary and obstetric factors. *Home Econ Res J* 1973;1:188-92.
28. Lechtig A, Yarbrough C, Delgado H, Habicht J, Martorell R, Klein R. Influence of maternal nutrition on birth weight. *Am J Clin Nutr* 1975;28:1223-33.
29. Turner RE, Langkamp HB, Littell RC, Lukowski MJ, Suarez MF. Comparing nutrient intake from food to the estimated average requirement shows middle to upper income pregnant women lack iron and possibly magnesium. *J Amer Dietetics Assoc* 2003;103:461-6.