

UNUSUAL PRESENTATION OF BENIGN CYSTIC TERATOMA

Chukwunwendu A. Okonkwo¹, Moses I. Momoh², Victor J. Ekanem³,
Maxwell E. Odiegwu⁴, Eugene M. Ikeanyi⁵, Ekene OMO⁶

ABSTRACT

Benign Cystic Teratoma (Dermoid cyst) is the most common Germ cell tumor, they rarely grow larger than 15cm in diameter and usually occur in young women with peak incidence between the ages of 20 and 40 years. We report a case of an unusually massive benign cystic teratoma measuring 86cm by 70cm and weighing 21kg, removed from a 58 year old six years post menopausal Para five woman who had been carrying the cyst over a period of twenty years. She had total abdominal hysterectomy and bilateral salpingo-oophorectomy and was discharged on the eighth post operative day.

KEY WORDS: Massive dermoid cyst, Postmenopausal woman.

Pak J Med Sci July - September 2008 Vol. 24 No. 4 624-626

How to cite this article:

Okonkwo CA, Momoh MI, Ekanem VJ, Odiegwu ME, Ikeanyi EM, Ekene OMO. Unusual presentation of benign cystic teratoma. Pak J Med Sci 2008;24(4):624-6.

INTRODUCTION

Germ cell tumors of the ovary form a group of neoplasm's comprised of many individual and mixed entities. Approximately 25-30% of all ovarian tumors are of Germ cell origin and of these, 95% are benign and only 3-4% are malignant.¹ Benign cystic teratomas (Dermoid cyst) account for 10-20% of all ovarian neoplasms.² They are encountered predominantly in women in their second and third decades of life and they rarely grow larger than 15cm.²

We report a case of an unusually massive benign cystic teratoma of the ovary, measuring 86cm by 70cm with a total weight of 21kg, (solid components weighing 12kg and cystic fluid 9litres) in a 58year old Para five lady who was six years post menopausal. She had carried this tumor for about twenty years.

CASE REPORT

Mrs. M. C. was a 58 year old Para five lady who presented to the Gynecology Clinic with a Twenty year history of an Abdominal mass

-
1. Dr. Chukwunwendu A. Okonkwo, MBBS, FMCOG, FICS
Department Of Obstetrics and Gynaecology
 2. Dr. Moses I. Momoh, MBBS, FWACS
Department of Surgery
 3. Dr. Victor J. Ekanem, MBBS, FMC PATH
Department of Morbid Anatomy
 4. Dr. Maxwell E. Odiegwu, MBBS, FMCOG
 5. Dr. Eugene M. Ikeanyi, MBBS
Department Of Obstetrics and Gynecology
 6. Dr. Ekene OMO, MBBS
Department of Anaesthesia
- 1-6: University of Benin Teaching Hospital,
Benin-city,
Edo State,
Nigeria.

Correspondence

Dr. Okonkwo CA,
Department of Obstetrics and Gynaecology,
University of Benin Teaching Hospital,
Benin-city,
Edo State,
Nigeria.

E-mail: drcaokonkwo@yahoo.com

* Received for Publication: February 8, 2008

* Accepted: June 1, 2008

which was increasing progressively in size. She had five spontaneous vaginal deliveries at home which were uneventful. She was six years post menopausal. She had tried unsuccessfully to obtain treatment through unorthodox medical practitioners for over 18 years and did not bother to seek proper medical attention because she could not afford it. A distant relative who was a medical practitioner undertook to pay her hospital bills.

The medical history and review of systems was unremarkable. Physical examination revealed a huge abdomino-pelvic mass with an abdominal girth of 120cm. An abdominal ultrasound demonstrated a huge cystic mass containing solid portions arising from the right ovary. Routine laboratory investigations done at admission were normal.

At exploratory laparotomy, a huge cyst occupying the entire abdomino-pelvic cavity with a smooth glistening surface was identified arising from the right ovary and attached by flimsy adhesions to the anterior aspect of the parietal peritoneum. The cyst had displaced bowel loops superiorly. The uterus, fallopian tubes and left ovary were normal. A total abdominal hysterectomy and bilateral salpingo-oophorectomy was done, commencing with oophorectomy of the massive right ovarian cyst and followed by total abdominal hysterectomy and salpingo-oophorectomy for the left ovary. The tumor was bisected post

operatively and was found to be cystic and filled with yellow sebum mixed with hair. The uterus and both ovaries were sent for pathological examination. The post operative period was uneventful and she was discharged on the eighth post operative day.

The histopathology report confirmed a dermoid cyst which showed matured tissues derived from the germ layers and consisting mostly of skin and its appendages, muscle and intestinal tissue.

DISCUSSION

Benign cystic teratomas, often called Ovarian Dermoids, account for 97% of ovarian teratomas and 10 - 20% of ovarian neoplasm's. They are usually around 5 - 15cm in diameter, often heavy for their size and frequently on a long pedicle. These latter features account for a ready liability to torsion.² The cyst can occur at any age, but 90% occur in women of reproductive age.² Mrs. M. C. was 58 years old and six years post menopausal. The cyst was 86cm by 70cm and weighed 21kg, despite its size, weight and long period of occurrence; the cyst did not undergo torsion.

This patient spent over 18 years seeking help for her situation. She visited several unorthodox medical practitioners. This type of health seeking behaviour is not uncommon amongst rural dwellers like Mrs. M. C, who lives in a village over 150km from the state capital,



Figure-1



Figure-2

where she presented. Most people in the rural community have a poor perception of hospitals. Some of their impressions are justified as most hospitals (private and government) are run incompetently; and hospital fees even for preventive measures as antenatal care, are usually arbitrary and unrealistic. Patients spend long hours at clinics and health workers are viewed as patient unfriendly.³

The surgical management of Benign cystic teratoma should be directed according to age, desire for further fertility and presence of concomitant pelvic pathology rather than size or the bilaterality status.⁴ Mrs. M .C. had completed her family and was post menopausal hence she had a total abdominal hysterectomy and bilateral salpingo-oophorectomy. She responded very well to the management and

was discharged on the eighth post operative day. This case highlights the very uncommon features dermoid cysts can present with and also shows the resultant effect on individual health due to socio-economic difficulties.

REFERENCES

1. Kurman RJ, Norris HJ. Malignant germ cell tumors of the ovary. Hum Pathol 1977;8:551-64.
2. Peel KR. Benign and Malignant tumors of the ovary; In Whitfield CR. Eds. Dewhursts textbook of Obstetrics and Gynaecology for Postgraduates. London, Blackwell Science Ltd.1995.
3. Okpere EE. Maternal Mortality in Nigeria. In: Okpere EE ed. Clinical Obstetrics: Benin-City; Uniben Press 2005;391-2.
4. Sah SP, Uprety D, Rani S. Germ cell tumors of the ovary: A clinicopathologic study of 121 cases from Nepal. J Obstet Gynaecol 2004;30(4):303-8.

Electronic Submission of Articles

“PAKISTAN JOURNAL OF MEDICAL SCIENCES” now accepts electronic submission of articles via e-mail, attachment in MS Word format at any of the following addresses:

pjms@pjms.com.pk
pulse@pulsepakistan.com

Note: The figures should be sent in JPEG format to ensure good quality images.

(Arrangements are also being made to accept manuscripts through our website in due course of time)

Electronic submission saves time, postage costs and allows the manuscript to be handled in electronic form throughout the publication process.

Processing Charges: Effective June 1st, 2008 all manuscripts from Pakistan should be accompanied with a Bank Draft for Rs. 500/- (Non-refundable) in the name of Pakistan Journal of Medical Sciences.

For detailed instructions to authors visit our website

pjms.com.pk