

Can blood and semen presepsin levels in males predict pregnancy in couples undergoing intra-cytoplasmic sperm injection?

Ali Ovayolu¹, Cansev Yilmaz Arslanbuğa², Ismet Gun³,
Belgin Devranoglu⁴, Arman Ozdemir⁵, Sule Eren Cakar⁶

ABSTRACT

Objective: To determine whether semen and plasma presepsin values measured in men with normozoospermia and oligoasthenospermia undergoing invitro-fertilization would be helpful in predicting ongoing pregnancy and live birth.

Methods: Group-I was defined as patients who had pregnancy after treatment and Group-II comprised those with no pregnancy. Semen and blood presepsin values were subsequently compared between the groups. Parametric comparisons were performed using Student's t-test, and non-parametric comparisons were conducted using the Mann-Whitney U test.

Results: There were 42 patients in Group-I and 72 in Group-II. In the context of successful pregnancy and live birth, semen presepsin values were statistically significantly higher in Group-I than in Group-II ($p=0.004$ and $p=0.037$, respectively). The most appropriate semen presepsin cut-off value for predicting both ongoing pregnancy and live birth was calculated as 199 pg/mL. Accordingly, their sensitivity was 64.5% to 59.3%, their specificity was 57.0% to 54.2%, and their positive predictive value was 37.0% to 29.6%, respectively; their negative predictive value was 80.4% in both instances.

Conclusion: Semen presepsin values could be a new marker that may enable the prediction of successful pregnancy and/or live birth. Its negative predictive values are especially high.

KEY WORDS: ICSI, Infertility, Oligoasthenoteratozoospermia, Presepsin, Seminal plasma.

doi: <http://dx.doi.org/10.12669/pjms.325.10610>

How to cite this:

Ovayolu A, Arslanbuğa CY, Gun I, Devranoglu B, Ozdemir A, Cakar SE. Can blood and semen presepsin levels in males predict pregnancy in couples undergoing intra-cytoplasmic sperm injection? *Pak J Med Sci.* 2016;32(5):1116-1120.

doi: <http://dx.doi.org/10.12669/pjms.325.10610>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Infertility is a frequently seen problem with a 10-15% rate of incidence, and is associated with male factors in as many as 50% of cases. A male infertility-associated factor is usually found together with abnormal semen parameters.¹ One of these is idiopathic oligoasthenoteratozoospermia (iOAT). iOAT unattributable to any specific cause accounts for about 24% of cases. OAT is diagnosed in accordance with the World Health Organization and/or Kruger strict criteria.^{2,3} Presepsin is a newly discovered substance that can be measured in blood and semen, and has been found to increase in many infections and non-infectious immune

Correspondence:

Ali Ovayolu,
Department of Obstetrics and Gynecology,
Cengiz Gokcek Public Hospital,
Gaziantep, Turkey,
Address: Osmangazi Mahallesi,
Cengiz Gökcek Kadın Hastalıkları ve Doğum,
Hastanesi 27010,
Gaziantep, Turkey.
E-mail: drovayolu@yahoo.com

- * Received for Publication: May 20, 2016
- * Revision Received: May 26, 2016
- * Edited and Corrected by Reviewer: July 26, 2016
- * Approved for Publication: * July 30, 2016

events.⁴⁻⁶ Cluster of differentiation 14 (CD14) is a glycoprotein expressed on the membrane surface of monocytes and macrophages and serves as a receptor for lipopolysaccharides (LPSs) and LPS-binding proteins (LPBs).

During inflammation, plasma protease activity generates soluble CD14 (sCD14) fragments. One of them, the sCD14 subtype (sCD14-ST), or presepsin, is normally present in very low concentrations in the serum of healthy individuals and has been shown to increase in response to bacterial infections.^{5,7,8} Procalcitonin and inflammatory mediators such as IL8 have been investigated in male infertility.⁹ Additionally, in many sexually-transmitted infections, interleukins were studied in systemic and local genital secretions.¹⁰

Focusing on couples with unexplained infertility and iOAT, our aim in this study was to research whether presepsin values of semen and blood plasma samples provided by the male partner on the oocyte pick-up day (OPU) could predict biochemical pregnancy, successful pregnancy, and live birth. No male patient with an infectious illness on the sample day was included in the study.

METHODS

This study was performed in the Department of Obstetrics and Gynecology, Zeynep Kamil Training and Educational Hospital, between July 2013 and July 2014, among infertile couples that were diagnosed as having iOAT or unexplained infertility (normozoospermia) for which no female-attributed cause could be found. The women were aged between 23 and 39 years; the patients were non-smokers and had a body mass index (BMI) <28 kg/m², and had no uterine, tubal or ovarian problems. Ovulation induction was performed using the classic long agonist protocol or fixed GnRH antagonist protocol. Both GnRH agonist and GnRH antagonist continued until the day of hCG administration to induce ovulation. On day 3 of cycle, gonadotropin treatment was initiated. The starting gonadotropin dose was individualized according to age, BMI, ovarian reserve determined by antral follicle count and basal FSH, and experience from previous cycles (between 150-300 IU). Serial ultrasonographic checks and estradiol (E2) level measurements were performed until 3 follicles were ≥17 mm and a serum E2 level >500 pg/mL were detected. Choriogonadotropin alpha 250 µg s.c. (Ovitrelle; MerckSerono, Turkey) was administered to induce final follicular maturation.

Oocyte retrieval took place 35.5-36 h after hCG administration. Intra-cytoplasmic sperm injection (ICSI) was performed after four hours of OPU. Fertilization was assessed at 16-18 h after ICSI. All embryo transfers were performed after OPU in two to five days.

According to national policy, the best morphologic grade one or two embryos were transferred into the uterine cavity under ultrasound guidance (GE logic alfa 200). Luteal support was initiated on the night of oocyte retrieval and continued until the day of pregnancy testing. If the test was positive, progesterone treatment was continued up to the 9th gestational week. Biochemical pregnancy was defined as a positive pregnancy test result (hCG levels > 20 mIU/mL) 12 days after embryo transfer. Patients that could not undergo embryo transfer were excluded from the study.

Seminal plasma samples were given for ICSI on the OPU day after 3 or 4 days of abstinence, and venous blood samples were taken and centrifuged at 2000 rpm for 10 minutes, and stored at 80°C until the required day for analysis. After determining pregnancy test results and live birth outcomes, samples were dissolved in a few hours and final measurements were performed. Patients were divided into two groups according to pregnancy occurrence. Group-I comprised those who were pregnant and Group-II was for those without pregnancy. Presepsin was measured using a PATHFAST chemiluminescence immunoassay analyzer (Mitsubishi Medience, Tokyo, Japan). Presepsin values measured and compared between the blood and semen groups.

Statistical analyses were performed using the Statistical Package for the Social Sciences for Windows version 15.0 software (SPSS, Chicago, IL., USA). Descriptive statistics are given as mean, standard deviation, frequency, and percentage. A parametric comparison was performed using Student's t-test, and a non-parametric comparison was conducted using the Mann-Whitney U test. Statistical significance was defined as $p < 0.05$.

Ethics Committee Approval: Approval was received for this study from the ethics committee of Zeynep Kamil Education and Research Hospital and written informed consent was obtained from patients who participated in this study.

RESULTS

In total, 114 patients were included in the study. There were 42 patients in Group-I and 72 patients

Table-I: Demographic, stimulation, and treatment outcome characteristics, and seminal plasma and plasma presepsin values of the groups.

Characteristics	Group I	Group II	P
Womens' age(years)	30.3±4.4	31.6±7.3	0.340*
Infertility duration, y	5.7±4.0	6.4±3.9	0.373*
FSH, mIU/ml	6.9±1.6	6.7±1.5	0.457*
Average used gonadotrophin, IU	2002.1±1001.8	2204.1±1033.8	0.325*
Mens' age(years)	33.5±4.8	34.3±7.6	0.548*
Biochemical pregnancy, n	n (42)	n (72)	
Semen presepsin, pg/mL	541.6±837.6	270.5±364.3	0.060 ^b
Plasma presepsin, pg/mL	343.2±217.3	358.6±206.1	0.458 ^b
Ongoing pregnancy, n	n (33)	n (81)	
Semen presepsin, pg/mL	644.1±927.1	261.2±346.1	0.004 ^b
Plasma presepsin, pg/mL	341.0±226.1	357.7±203.6	0.420 ^b
Live birth, n	n (28)	n (86)	
Semen presepsin, pg/mL	643.0±973.2	280.0±367.7	0.037 ^b
Plasma presepsin, pg/mL	343.2±225.4	356.2±205.1	0.521 ^b

Data are presented as mean ± SD. *Student's t-test and bMann-Whitney U test.

in Group-II. Table-I indicates that there was no statistically significant difference between the groups in terms of demographic and treatment characteristics. The rate of ongoing pregnancy was 78.6%, and the rate of live birth was 66.7%. A comparative examination of the semen and plasma presepsin values revealed higher values in Group-I than in the non-pregnant Group-II in terms of biochemical pregnancy, successful pregnancy, and live birth. However, in terms of successful pregnancy and live birth, only semen presepsin values were statistically significantly higher in Group I than in Group II ($p=0.004$ and $p=0.037$, respectively). Table-II shows that there is no statistical difference

between the oligoasthenospermic group and the normozoospermic group in terms of semen and plasma presepsin values.

Fig.1 shows the ROC curve, which indicates the efficiency of semen presepsin values in predicting successful pregnancy. Fig.2 indicates the efficiency of semen presepsin values in predicting live birth. In the context of successful pregnancy, the area under the curve (AUC) for semen presepsin values was 0.677, 95% confidence interval (CI):[0.563-0.792]; $p=0.004$. The most appropriate semen presepsin cut-off point in predicting successful pregnancy was 199 pg/mL. Using 199 pg/mL as




Fig.1




Fig.2

Table-II: Seminal plasma and plasma presepsin values in the oligoasthenospermia and normozoospermia groups.

Characteristics	Oligoasthenospermia n (61)	Normozoospermia n (53)	P
Mens' age, years	33.3±5.3	34.9±8.0	0.182*
Seminal plasma Presepsin, pg/mL	371.6±490.4	366.3±695.75	0.829b
Plasma Presepsin, pg/mL	377.6±233.0	324.3±176.6	0.338b

Data are presented as mean ± SD. *Student's t-test and bMann-Whitney U test.

the cut-off point, the sensitivity and specificity of presepsin values in predicting successful pregnancy were 64.5% and 57.0%, and their positive and negative predictive values were 37.0% and 80.4%, respectively. In the context of live birth, the AUC for semen presepsin values was 0.634, 95% CI: [0.509-0.759]; $p=0.037$. The most appropriate semen presepsin breakpoint in predicting live birth was again 199 pg/mL. Using 199 pg/mL as the cut-off point, the sensitivity and specificity of presepsin values in predicting live birth was 59.3% and 54.2%, and their positive and negative predictive values were 29.6% and 80.4%, respectively.

DISCUSSION

When the studies related to presepsin are analyzed, it is clear that it represents acute inflammatory events.⁶ In this study we did not measure "fresh" presepsin immediately. We think that this does not pose a problem when compared with previous studies; however, the number of similar studies is not large and perhaps this approach is not the most accurate.^{11,12} Although the diagnostic significance of serum presepsin measurements have been established, no reports have addressed the presence of this peptide in seminal fluid. We observed that it was possible to measure presepsin in semen and that patients can have different values. We also observed that the same patient could have different seminal plasma and plasma presepsin values. The main aim of the study was to find out a new marker that could predict successful pregnancy and/or live birth in infertile patient groups. Our ROC curve results indicate that the AUC values both in the context of successful pregnancy and live birth were 0.677 and 0.634, respectively.

As is known, in order to be a good indicator, AUC values have to be 0.80 and above. However, the small sample size in our study may have prevented us attaining such a level. As the variants in our study were not homogenous, we had to use a non-parametric test. In addition, standard deviation of semen and plasma presepsin values

were found to be very high. This situation could correspond to a reconsideration of presepsin measurement standards. Nonetheless, presepsin values were found to be statistically significant in both successful pregnancy and live birth ($p=0.004$ and $p=0.037$, respectively). As a result of the ROC curve analysis, the cut-off point most appropriate for both positions was calculated as 199 pg/mL. When semen presepsin values were taken to be 199 pg/mL and above as a diagnostic value in predicting successful pregnancy and live birth, the negative predictive values were high and at a good level, 80.4%.

In conclusion, semen presepsin values could be a new diagnostic test for predicting ongoing pregnancy and/or live birth. In particular, their negative predictive value is quite strong. This preliminary study is weak in numeric terms. Further studies with larger sample sizes are necessary to confirm our preliminary findings.

ACKNOWLEDGMENTS

We would like to thank our dear colleague Dr. Selim Kilic, a member of the Clinic Department of Epidemiology and Public Health, Ankara Gülhane Military Medical School for helping with the statistical analysis of this study. We also thank our colleagues Dr Tayfun Kutlu, Dr Naciye Arat, and Dr Gülden Tunali from the IVF unit of Zeynep Kamil Education and Research Hospital for providing insight, expertise, and scientific guidance that greatly assisted our research.

Declaration of Interest: In this study Presepsin was sponsored by Biobak commercial company (Turkey).

Financial Disclosure: The authors declared that this study has received no financial support.

Note: This manuscript was presented as a poster at Vth Society of Reproductive Medicine and Surgery Congress between 28th - 31st October, 2015 at Cornelia Diamond Hotel, Belek, Antalya.

REFERENCES

1. Bhasin S, de Kretser DM, Baker HW. Clinical review 64: Pathophysiology and natural history of male infertility. *J Clin Endocrinol Metab*. 1994;79:1525-1529. doi: 10.1210/jcem.79.6.7989450
2. World Health Organization. WHO laboratory manual for the examination and processing of human semen. 5th ed. Geneva: World Health Organization. 2010; pp. 271. Available from: http://apps.who.int/iris/bitstream/10665/44261/1/9789241547789_eng.pdf
3. Kruger TF, Acosta AA, Simmons KF, Swanson RJ, Matta JF, Oehninger S. Predictive value of abnormal sperm morphology in in vitro fertilization. *Fertil Steril*. 1988;49:112-127.
4. Shirakawa K, Naitou K, Hirose J, Takahashi T, Furusako S. Presepsin (sCD14-ST): development and evaluation of one-step ELISA with a new standard that is similar to the form of presepsin in septic patients. *Clin Chem Lab Med*. 2011;49(5):937-939. doi: 10.1515/CCLM.2011.145.
5. Endo S, Suzuki Y, Takahashi G, Shozushima T, Ishikura H, Murai A, et al. Presepsin as a powerful monitoring tool for the prognosis and treatment of sepsis: a multicenter prospective study. *J Infect Chemother*. 2014;20:30-34. doi: 10.1016/j.jiac.2013.07.005.
6. Zou Q, Wen W, Zhang XC. Presepsin as a novel sepsis biomarker. *World J Emerg Med*. 2014;5(1):16-19. doi: 10.5847/wjem.j.1920-8642.2014.01.002.
7. Mussap M, Noto A, Fravega M, Fanos V. Soluble CD14 subtype presepsin (sCD14-ST) and lipopolysaccharide binding protein (LBP) in neonatal sepsis: new clinical and analytical perspectives for two old biomarkers. *J Matern Fetal Neonatal Med*. 2011;24:12-14. doi: 10.3109/14767058.2011.601923.
8. Yaegashi Y, Shirakawa K, Sato N, Suzuki Y, Kojika M, Imai S, et al. Evaluation of a newly identified soluble CD14 subtype as a marker for sepsis. *J Infect Chemother*. 2005;11:234-238. doi: 10.1007/s10156-005-0400-4
9. Slavakis A, Papadimas J. Procalcitonin: does it play a role in male reproduction? *Fertil Steril*. 2000;74(6):1227-1228. doi: 10.1016/S0015-0282(00)01588-0
10. Hedges SR, Sibley DA, Mayo MS, Hook EW 3rd, Russel MW. Cytokine and antibody responses in women infected with *Neisseria gonorrhoeae* effects of concomitant infections. *J Infect Dis*. 1998;178(3):742-751.
11. Okamura Y, Yokoi H. Development of a point-of-care assay system for measurement of presepsin (sCD14-ST). *Clin Chim Acta*. 2011;412:2157-2161. doi: 10.1016/j.cca.2011.07.024
12. Wu J, Hu L, Zhang G, Wu F, He T. Accuracy of presepsin in sepsis diagnosis: A systematic review and meta-analysis. *PLoS ONE*. 2015;10(7):e0133057. doi: 10.1371/journal.pone.0133057

Authors' Contribution:

AO, CYA, IG conceived, designed and did statistical analysis & editing of manuscript.

AO, CYA, AO, BD, SEC did data collection and manuscript writing.

AO, IG, BD, AO, SEC did review and final approval of manuscript.

Authors:

1. Dr. Ali Ovayolu,
Department of Obstetrics and Gynecology,
Cengiz Gokcek Public Hospital,
Gaziantep, Turkey.
2. Dr. Cansev Yilmaz Arslanbuga,
Department of Clinical Microbiology,
3. Dr. Ismet Gun,
Associate Professor,
Department of Obstetrics and Gynecology,
GATA Faculty of Medicine,
Istanbul, Turkey.
4. Dr. Belgin Devranoglu,
Department of Obstetrics and Gynecology,
5. Dr. Arman Ozdemir,
Department of Urology,
6. Dr. Sule Eren Cakar,
Department of Clinical Microbiology,
- 2,4-6: Zeynep Kamil Education and Research Hospital,
Istanbul, Turkey.