

Correspondence

Use of Guedel Airway as a guide to insert nasogastric tube under general anaesthesia: A simple and logical way

Faraz Shafiq¹, Faruq Hameed², Khalid Siddiqui³

KEYWORDS: Guedel, Nasogastric, Anaesthesia.

doi: <https://doi.org/10.12669/pjms.345.16572>

How to cite this:

Shafiq F, Hameed F, Siddiqui K. Use of Guedel Airway as a guide to insert nasogastric tube under general anaesthesia: A simple and logical way. *Pak J Med Sci*. 2018;34(5):1305-1306. doi: <https://doi.org/10.12669/pjms.345.16572>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dear Editor,

We all must have faced difficulty in inserting nasogastric tube (NGT) during general anaesthesia (GA). Procedure seems to be very straightforward but has got consequences in terms of failure, which can cause bleeding and alteration in hemodynamic variables. The overall failure rate turned out to be around 50-60%.¹ The kinking or coiling of distal part of tube in piriform sinus or at level of arytenoids cartilage² is reported to be the frequent cause of this failure. However, it's still not very clear, as many factors including anatomical variations of oropharynx or type of material used by manufacturer may be responsible. We hypothesized that base of tongue and oropharynx is the possible site of resistance causing difficulty to insert NGT. The situation get worsen in patients having GA due to relaxation of tongue and jaw

muscles.³ (Fig.1). The use of proper size Guedel oro-pharyngeal airway may be beneficial for NGT insertion in patients under GA. (Fig.2). It does




Fig.1: Loss of tone at back of tongue during GA causing resistance to NGT passage.




Fig.2: Presence of Guedel airway prevents tongue fall and assisting NGT to pass down.

1. Faraz Shafiq,
Assistant Professor,
2. Faruq Hameed,
Resident,
3. Khalid Siddiqui,
Assistant Professor,
- 1-3: Department of Anaesthesiology,
The Aga Khan University,
Karachi Pakistan.

Correspondence:

Faraz Shafiq
Assistant Professor,
Department of Anaesthesiology,
The Aga Khan University,
Karachi, 74800, Pakistan.
Email: farazshafiq76@yahoo.com

* Received for Publication: August 31, 2018
* Accepted for Publication: September 26, 2018

this by preventing tongue fall⁴ minimizing the resistance and likelihood of NGT placement in first pass. We found this technique really useful. However recommend further clinical trial to judge its success alone or as at adjuvant maneuvers for NGT placement.

Acknowledgments: We acknowledge the efforts made by *Mr. Aslam Bashir* in drawing medical illustrations for this report.

Conflict of interest: None.

REFERENCES

1. Appukutty J, Shroff PP: Nasogastric tube insertion using different techniques in anesthetized patients: a prospective, randomized study. *Anesth Analg.* 2009;109:832-835.
2. Parris WC. Reverse Sellick maneuver. *Anesth Analg.* 1989;68:423.
3. Boidin MP. Airway patency in the unconscious patient. *Br J Anaesth.* 1985;57:306-310.
4. Baskett TF. Arthur Guedel and the oropharyngeal airway. *Resuscitation.* 2004;63:3-5.