

Assessment of adipokines relationships with cardiovascular risk markers in relation to body indices in normoglycemic males

Syed Shahid Habib¹, Khilid A Al Regaeiy², Laila Al Dokhi³

ABSTRACT

Objectives: To evaluate the phenotypic relationship between obesity indices, resistin, adiponectin and cardiovascular risk markers in normoglycemic healthy individuals.

Methodology: This cross-sectional study was conducted in the Department of Physiology College of Medicine, King Saud University, Riyadh. A total of 120 male subjects were selected for the study. All subjects underwent analysis of body composition, glucose, glycosylated hemoglobin (HbA1c), lipids, adiponectin, resistin, lipoprotein(a) and high sensitivity C reactive protein (hsCRP).

Results: Body mass index (BMI) ($r=0.326$, $p < 0.001$), body fat mass (BFM) ($r=0.377$, $p < 0.001$), body fat percentage (BF%) ($r=0.326$, $p < 0.001$), waist hip ratio (WHR) ($r=0.402$, $p < 0.001$) and basal Insulin levels ($r=0.217$, $p=0.018$) were positively correlated with hsCRP. However, serum adiponectin levels ($r=0.189$, $p=0.0391$) were negatively correlated with hsCRP. Adiponectin levels were significantly lower in obese compared to non obese subjects ($p=0.0551$). Keeping hsCRP as dependant variable we observed that WHR, BFM, BF%, BMI and adiponectin were significant predictors in univariate analysis. In multiple regression analysis WHR and adiponectin were independent predictors of hsCRP.

Conclusion: Obese individuals have significantly higher levels of hsCRP levels and lower adiponectin levels than non obese subjects. Serum adiponectin levels and WHR are independant predictors of hsCRP levels in normoglycemic subjects.

KEY WORDS: Body composition, Fat mass, Body fat percentage, High sensitivity C reactive protein, Lipoprotein(a).

doi: <http://dx.doi.org/10.12669/pjms.291.2913>

How to cite this:

Habib SS, Al-Regaeiy KA, Al-Dokhi A. Assessment of adipokines relationships with cardiovascular risk markers in relation to body indices in normoglycemic males. *Pak J Med Sci* 2013;29(1):21-26. doi: <http://dx.doi.org/10.12669/pjms.291.2913>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Syed Shahid Habib, FCPS,
2. Khilid A Al Regaeiy, PhD,
3. Laila Al Dokhi, PhD,
- 1-3: Department of Physiology, College of Medicine, King Saud University, Riyadh-11461, Kingdom of Saudi Arabia.

Correspondence:

Dr. Syed Shahid Habib,
Associate Professor,
Department of Physiology (29),
College of Medicine,
PO Box 2925,
King Saud University, Riyadh-11461,
Kingdom of Saudi Arabia.
E-mail: shahidhabib44@hotmail.com

* Received for Publication: September 10, 2012

* Accepted for Publication: October 11, 2012

INTRODUCTION

It is apparent that obesity represents one of the foremost contributory factors leading to diabetes.¹ Metabolic syndrome is associated with high risk for the development of diabetes and cardiovascular diseases (CVD).² The greater risk of type 2 Diabetes Mellitus (DM) in the obese can, at least partly, be explained by changes in adipose tissue function.³⁻⁵ It is now well known that adipose tissues are active endocrine tissues that produce >50 adipokines, including leptin, adiponectin, resistin, interleukin-6 (IL-6), tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), vascular endothelial growth factors, nerve growth factors, and haptoglobin.⁶

The recent interests in adipocyte derived factors have resulted in identification of a large group of adipocyte specific proteins in blood, such as adiponectin, acylation stimulating proteins, resistin and leptin.⁷ These adipocyte derived hormones are presently under intensive investigations concerning their involvement in the regulation of adipose tissue physiology, and in particular, their potential implication in insulin resistance, obesity and diabetes.⁸ Resistin is a protein hormone secreted by adipocytes which leads to insulin resistance and is considered to be an important link between obesity and diabetes.⁹ Resistin has been involved in the pathogenesis of obesity-mediated insulin resistance and type 2 DM. The expression and functional properties of adipocytokines and their effects on metabolism have been under intensive investigation. Indeed, studies on adipocytokines and their potential effects in human obesity and type 2 DM have implicated them in the pathogenesis of metabolic syndrome.¹⁰ However, still the true pathogenic mechanism that leads to accelerated atherogenesis in DM is still not known, because of its multi system ramifications. Elevated Lipoprotein(a) [Lp(a)] levels confer genetic predisposition to CVD and may be one of the links to accelerated atherogenesis in DM.¹¹⁻¹³ Our previous reports have shown that Lp(a) is a marker of the presence, diffuseness and severity of CAD in Saudi adults.¹⁴

A large body of evidence have shown that the inflammatory biomarker high-sensitivity C-reactive protein (hsCRP) is an independent predictor of cardiovascular events and it also predicts risk of incident hypertension and diabetes.¹⁵ Although it is apparent that inconsistencies remain in the data for a role of adipokines in obesity, there is a compelling evidence for their involvement in the etiology of insulin resistance and type 2 DM. The relationship among obesity, inflammatory markers (such as adipokines, phase reactant proteins), and insulin resistance (IR) has been investigated extensively in several populations but not explicitly in Saudi arabian population in which the prevalence of obesity and associated comorbidities is one of the highest in the world. Men have considerably lower plasma adiponectin levels than women, concordant with previous studies.¹⁶ Therefore we studied only one gender to exclude the possible confounding effect of gender on variables studied. This may partly relate to inhibitory effects of androgens on adiponectin levels.¹⁷

To the best of our knowledge there are few studies relating both resistin and adiponectin

in normoglycemic subjects with obesity indices. Therefore, this project evaluated the phenotypic relationship between obesity indices, resistin, adiponectin and cardiovascular risk markers in normoglycemic healthy individuals.

METHODOLOGY

Subjects: This cross-sectional study was conducted in the Department of Physiology College of Medicine, King Saud University, Riyadh. Patients were recruited from King Khalid University Hospital. The consent form was designed both in Arabic and English and was approved by the College of Medicine ethics review board. A total of 120 male subjects were selected for the study. All subjects underwent Body composition analysis.

Laboratory Analysis: Overnight fasting blood samples were collected, and analyzed for fasting blood glucose (FBG), total cholesterol (TC), Triglycerides (TG), Low density Lipoprotein (LDL), High density lipoprotein (HDL), adiponectin, resistin, Lp(a) and hsCRP. Human hsCRP and Lp(a) immunoassays were performed by quantitative standard sandwich ELISA technique using monoclonal antibody specific for these analytes with kits supplied by IBL International GMBH Germany. The results of patients with hsCRP values >10 mg/L were discarded and were re-evaluated after 2-3 weeks. We followed the guidelines of the American Heart Association for measurement, evaluation and expression of hsCRP.¹⁸

Statistical Analysis: The data was analyzed by computer software program Statistical Package for Social Sciences (SPSS Version 19, Chicago, IL). Descriptive characteristics of the study patients were calculated as mean±SD (Standard Deviation) for continuous variables. The tests applied for statistical analysis were Student's t test for normally distributed and non-parametric Mann-Whitney U test for data which was not following normal distribution. Univariate correlations of Serum hsCRP with body indices, fat mass and other metabolic parameters were evaluated with both Spearman's rank order and Pearson correlations. Stepwise multiple regression analyses were performed to further identify the most significant variables contributing to hsCRP levels. A p value of ≤0.05 was taken as statistically significant.

RESULTS

Anthropometric and biochemical profiles of participants included in this study are displayed in

Table-I: Anthropometric and metabolic characteristics of the subjects and correlation coefficients with hsCRP.

Total Subjects N=120	Mean±SD	Minimum	Maximum	r	p value
Age years	40.16±11.81	21.00	71.00	0.186	0.044
Height cm	167.42±8.21	147.00	187.00	-0.174	0.059
Weight kg	78.85±14.17	45.80	125.00	0.160	0.084
BMI kg/m ²	28.13±4.80	15.60	42.30	0.326	<0.001
Fat Mass kg	24.08±8.58	6.20	52.70	0.351	<0.001
% FAT	30.01±7.58	12.50	49.80	0.377	<0.001
WHR	0.95±0.07	0.77	1.16	0.402	<0.001
FBS mmol/l	5.04±0.91	2.90	5.60	-0.047	0.617
Insulin uIU/ml	24.20±11.87	14.15	132.15	0.217	0.018
HOMA-IR	5.50±3.35	2.58	32.30	0.130	0.159
HbA1c %	5.01±0.60	2.90	5.91	-0.021	0.833
TG mmol/l	1.14±0.72	1.29	6.20	0.115	0.281
TC mmol/l	4.64±0.91	2.90	6.20	0.119	0.267
HDL mmol/l	1.19±0.22	0.75	1.63	-0.073	0.504
LDL mmol/l	2.83±0.79	1.61	4.48	0.029	0.786
Adiponectin ng/ml	127.42±58.28	36.72	308.25	-0.115	0.216
Resistin ng/ml	2.40±1.14	0.57	5.58	0.088	0.342
Lp(a) mg/dl	24.45±20.55	2.00	90.04	0.106	0.252
hsCRP mg/L	3.75±2.36	0.81	9.80		

Data are expressed as Mean±SD with ranges. BMI, body mass index; % Fat, percent fat mass; HOMA-IR, homeostasis model assessment of insulin resistance; FBG, fasting blood glucose; TC, total cholesterol; TG, Triglycerides; LDL, Low density lipoprotein; HDL, High density lipoprotein; Lp(a), Lipoprotein(a); high-sensitivity C-reactive protein (hsCRP)

Table-I with correlation coefficients with hsCRP. We observed that BMI ($r=0.326$, $p<0.001$), BFM ($r=0.377$, $p<0.001$), BF% ($r=0.326$, $p<0.001$), WHR ($r=0.402$, $p<0.001$), Insulin ($r=0.217$, $p=0.018$) were positively correlated with hsCRP. However, serum adiponectin levels ($r=0.189$, $p=0.0391$) were negatively correlated with hsCRP.

Table-II shows comparison of Anthropometric and biochemical profiles, adipokines and CV risk markers between obese and non obese subjects. It was observed that obese subjects had significantly higher FBS ($p=0.0466$), HOMA-IR ($p=0.0002$), HbA1c ($p=0.0366$), TG ($p=0.0198$), hsCRP ($p=0.0096$) levels. While adiponectin levels were significantly lower in obese subjects compared to non obese subjects ($p=0.0551$).

In univariate regression analysis keeping hsCRP as dependant variable we observed that WHR, BFM, BF%, BMI and adiponectin were significant predictors. However, no correlation was observed with resistin. (Fig.1)

The degree to which obesity markers independently predicted hsCRP was examined in a series of multiple regression models. In these models, hsCRP was the dependent variable and obesity markers, adipokines, HOMAIR and lipids were the independent variables. In multiple regression analysis many variables became non significant predictors. Only WHR and adiponectin levels turned out to be independent predictors of hsCRP and this relationship was independent of age. (Table-III)

DISCUSSION

We evaluated the relationship between obesity indices (BMI, WHR, BF%, BFM), hsCRP, adipokines and IR in Saudi population. To the best of our knowledge, this study is the first to compare the

Table-II: Comparison of anthropometric and metabolic characteristics of between obese and non obese subjects.

	Non Obese Mean ±SD	Obese	P value
Age	40.18±10.99	40.12±13.34	0.9788
Height	167±7.56	166.40±9.28	0.3224
Weight	72.53±11.18	90.28±11.68	0.0000
BMI kg/m ²	25.56±3.40	32.85±2.99	0.0000
Fat Mass kg	19.63±5.71	32.11±6.89	0.0000
% FAT	26.82±6.39	35.69±6.06	0.0000
WHR	0.92±0.05	1.01±0.06	0.0000
FBS mmol/l	4.92±0.57	5.27±1.30	0.0446
Insulin uIU/ml	23.28±13.82	25.91±6.78	0.2485
HOMA-IR	4.75±1.40	5.97±1.99	0.0002
HbA1c	4.94±0.57	5.20±0.59	0.0366
TG mmol/l	1.06±0.51	1.37±0.73	0.0198
TC mmol/l	4.64±0.93	4.64±0.91	0.9757
HDL mmol/l	1.06±0.51	1.30±0.23	0.6933
LDL mmol/l	2.83±0.80	2.81±0.80	0.9017
Adiponectin ng/ml	130.04±54.91	112.53±42.84	0.0451
Resistin ng/ml	2.42±1.15	2.40±1.13	0.9351
Lp(a) mg/dl	24.79±21.95	23.85±17.98	0.8133
hsCRP mg/L	3.33±2.00	4.48±2.74	0.0096




Fig.1: Linear regression analysis of hsCRP (mg/L) as dependant variable (Y axis) with body indices and adipokines as independent variables (X axis).

associations among obesity, inflammation, adipokines and IR in Saudis. Results from this study confirmed previously reported associations between markers of obesity (BMI and WHR) and IR measured by HOMA.¹⁹⁻²¹

This study reports that serum adiponectin concentrations were inversely related to circulating hsCRP. Our data also confirms that BF% and BFM obesity are major determinants of plasma adiponectin concentrations. Higher levels of BF%, BFM and WHR, reflecting total and visceral

adiposity, were associated with an additive fall in plasma adiponectin concentrations. Similar to our reports Matsushita et al.²² performed a cross-sectional study of 624 Japanese middle-aged men and found that adiponectin level was a more significant predictor compared to TNF α , IL-6 or hsCRP for prevalent metabolic syndrome. In line with our results previous studies have also shown that obesity, especially visceral obesity, is associated with low-grade inflammation, which is characterized by an increase in plasma concentration

Table-III: Multiple regression analysis for prediction of hsCRP in 120 healthy middle aged men.

	Standardized Coefficients Beta	P value	95.0% Confidence Interval for B (Lower-Upper)
Age years	-.019	.897	-.064-.057
Fat Mass kg	.575	.252	-.132-.492
Weight kg	-.688	.329	-.370-.126
Height cm	.254	.580	-.196-.347
BF%	-.218	.579	-.340-.192
WHR	.669	.011	5.474-41.305
BMI kg/m ²	-.057	.913	-.627-.562
FBS mmol/dl	.410	.370	-1.941-5.138
Insulin uIU/ml	1.110	.123	-.127-1.036
HOMA-IR	-1.110	.161	-4.529-.772
HbA1c%	.026	.930	-2.202-2.406
TG mmol/L	-.038	.799	-1.087-.841
TC mmol/L	.236	.362	-.761-2.054
HDL mmol/L	-.039	.748	-3.154-2.278
LDL mmol/L	-.444	.070	-2.952-.117
Adiponectin ng/ml	-.251	.049	-.021- -.001
Resistin ng/ml	.067	.612	-.409-.688
Lp(a) mg/dl	.150	.211	-.010-.043

significant association among hsCRP, BMI, WHR, BF% & BFM in Saudis.²⁴⁻²⁶ Unlike most other adipocytederived hormones, circulating adiponectin levels are decreased in obesity.^{27,28} Adiponectin increases fatty acid oxidation and directly sensitizes the body to insulin.^{29,30}

The degree to which hsCRP may be predicted by markers of obesity were also assessed in different univariate and multivariate regression models. Taken together, our model showed that adiponectin and WHR were independent predictors of hsCRP for CV risk. Following the above observation, it is reasonable to suggest that the relationship between adipokines and hsCRP is to a significant extent dependent on visceral adiposity especially WHR.

Potential limitation of our study is its cross-sectional design from which we are not able to provide a strong causative conclusion. Secondly, our subjects were relatively homogenous with lower CVD risks because all were normoglycemic. Although various confounding factors such as sex, age and cigarette smoking have been minimized, we should be cautious to apply the conclusion to general population. Therefore, further extending the study samples to women and other ethnic populations with various CVD risks is needed.

CONCLUSION

Obese individuals have significantly higher levels of hsCRP levels and lower adiponectin levels than non obese subjects. Serum adiponectin levels and WHR are independent predictors of hsCRP levels in normoglycemic subjects.

ACKNOWLEDGMENTS

The authors are thankful to Mr. James Chu for blood assays and Mr. Timhar Amlih for collection of data. This work was supported by grant from King Abdul Aziz City for Science and Technology (Grant No: 26-60).

Conflict of interest: The authors have no conflicting interests and is or not supported/funded by any Drug Company.

REFERENCES

- Jensen MD. Role of body fat distribution and the metabolic complications of obesity. *J Clin Endocrinol Metab.* 2008;93:S57-S63.
- Ford ES. Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: a summary of the evidence. *Diabetes Care.* 2005;28:1769-1778.
- Bastard JP, Maachi M, Lagathu C, Kim MJ, Caron M, Vidal H, et al. Recent advances in the relationship between obesity, inflammation, and insulin resistance. *Eur Cytokine Netw.* 2006;17(1):4-12.
- Hajer GR, van Haeften TW, Visseren FLJ. Adipose tissue dysfunction in obesity, diabetes, and vascular diseases. *Eur Heart J.* 2008;29:2959-2971.
- Rasouli N, Kern PA. Adipocytokines and the metabolic complications of obesity. *J Clin Endocrinol Metab.* 2008;93(11 Suppl 1):S64-S73.
- Trayhurn P, Wood IS. Signaling role of adipose tissue: adipokines and inflammation in obesity. *Biochem Soc Trans.* 2005;33:1078-1081
- Faraj M, Lu HL, Cianflone K. Diabetes, lipids, and adipocyte secretagogues. *Biochem Cell Biol.* 2004;82:170-190.
- Silha JV, Krsek M, Skrha JV, Sucharda P, Nyomba BL, Murphy LJ. Plasma resistin, adiponectin and leptin levels in lean and obese subjects: correlations with insulin resistance. *Eur J Endocrinol.* 2003;149:331-335.
- Flier JS. Diabetes. The missing link with obesity? *Nature.* 2001;409:292-293.
- Kusminski CM, McTernan PG, Kumar S. Role of resistin in obesity, insulin resistance and Type II diabetes. *Clinical Sci.* 2005;109(3):243-256.
- Gambhir JK, Kaur H, Prabhu KM, Morrisett JD, Gambhir DS. Association between lipoprotein(a) levels, apo(a) isoforms and family history of premature CAD in young Asian Indians. *Clin Biochem.* 2008;41(7-8):453-458.
- de Brouckère V, Liénart F, Ducobu J. New data about lipoprotein(a): a curious molecule or a causal cardiovascular risk factor? *Rev Med Brux.* 2010;31(3):171-176.
- Anuurad E, Enkhmaa B, Berglund L. Enigmatic role of lipoprotein(a) in cardiovascular disease. *Clin Transl Sci.* 2010;3(6):327-332.

14. Habib SS, Abdel-Gader AM, Kurdi MI, Al-Aseri Z, Soliman MM. Lipoproteina(a) is a feature of the presence, diffuseness, and severity of coronary artery disease in Saudi population. *Saudi Med J*. 2009;30(3):346-352.
15. Ridker PM. C-reactive protein and the prediction of cardiovascular events among those at intermediate risk: moving an inflammatory hypothesis toward consensus. *J Am Coll Cardiol*. 2007;49(21):2129-2138.
16. Hung J, McQuillan BM, Thompson PL, Beilby JP. Circulating adiponectin levels associate with inflammatory markers, insulin resistance and metabolic syndrome independent of obesity. *Int J Obesity* 2008;32:772-779.
17. Nishizawa H, Shimomura I, Kishida K, Maeda N, Kuriyama H, Nagaretani H, et al. Androgens decrease plasma adiponectin, an insulin-sensitizing adipocyte-derived protein. *Diabetes*. 2002;51(9):2734-2741.
18. Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO, Criqui M, et al. Markers of inflammation and cardiovascular disease: application to clinical and public health practice. *Circulation*. 2003;107(3):499-511.
19. Leinonen E, Hurt-Camejo E, Wiklund O, Hultén LM, Hiukka A, Taskinen MR. Insulin resistance and adiposity correlate with acute-phase reaction and soluble cell adhesion molecules in type 2 diabetes. *Atherosclerosis*. 2003;166:387-394.
20. Hayashi T, Boyko EJ, McNeely MJ, Leonetti DL, Kahn SE, Fujimoto WY. Visceral adiposity, not abdominal subcutaneous fat area, is associated with an increase in future insulin resistance in Japanese Americans. *Diabetes*. 2008;57:1269-1275.
21. Garg A. Regional adiposity and insulin resistance. *J Clin Endocrinol Metab*. 2004;89(9):4206-4210.
22. Matsushita K, Yatsuya H, Tamakoshi K, Wada K, Otsuka R, Takefuji S, et al. Comparison of circulating adiponectin and proinflammatory markers regarding their association with metabolic syndrome in Japanese men. *Arterioscler Thromb Vasc Biol*. 2006;26:871-876.
23. Wellen KE, Hotamisligil GS. Obesity-induced inflammatory changes in adipose tissue. *J Clin Invest*. 2003;112(12):1785-1788.
24. Saijo Y, Kiyota N, Kawasaki Y, Miyazaki Y, Kashimura J, Fukuda M, et al. Relationship between C-reactive protein and visceral adipose tissue in healthy Japanese subjects. *Diabetes Obes Metab*. 2004;6:249-258.
25. Greenfield JR, Samaras K, Jenkins AB, Kelly PJ, Spector TD, Gallimore JR, et al. Obesity is an important determinant of baseline serum C-reactive protein concentration in monozygotic twins, independent of genetic influences. *Circulation*. 2004;109:3022-3028.
26. Doumatey AP, Lashley KS, Huang H, Zhou J, Chen G, Amoah A, et al. Relationships Among Obesity, Inflammation, and Insulin Resistance in African Americans and West Africans. *Obesity*. 2010;18:598-603.
27. Hu E, Liang P, Spiegelman BM. AdipoQ is a novel adipose-specific gene dysregulated in obesity. *J Biol Chem*. 1996;271:10697-10703.
28. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun*. 1999;257:79-83.
29. Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, et al. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating amp-activated protein kinase. *Nat Med*. 2002;8:1288-1295.
30. Berg AH, Combs TP, Du X, Brownlee M, Scherer PE. The adipocyte-secreted protein acrp30 enhances hepatic insulin action. *Nat Med*. 2001;7(8):947-953.