

Clinical characteristics of *Clostridium difficile*-associated diarrhea among patients in a tertiary care center in China

Yongqiang Li¹, Yi Huang², Yuyuan Li³, Yuqiang Nie⁴

ABSTRACT

Objective: This study investigated the incidence, risk factors, and clinical characteristics of *Clostridium difficile*-associated diarrhea (CDAD) in Chinese patients.

Methods: Fecal specimens of patients with antibiotic-associated diarrhea (AAD) were collected to test *C. difficile* toxin A and B using enzyme-linked fluorescent assay to identify CDAD. By adopting a nested case-control design, the matched people (ratio 1:3) without AAD were included as controls.

Results: Out of 56,172 inpatients, 39,882 (71.0%) used antibiotics, 470 suffered from AAD, and 93 were diagnosed with CDAD. The incidence of nosocomial CDAD was 166 per 100,000. The proportion of CDAD in AAD was 19.8%. CDAD patients presented with more severe clinical manifestations and exhibited more concurrent illness. Logistic regression analysis showed the risk factors of CDAD: advanced age, nasogastric tube-feeding, high APACHE II scores, high level of serum C-reaction protein, low level of serum albumin, severe underlying disease or comorbidity, and number of antibiotic intake. Twenty-nine patients (31.2%) were cured with vancomycin, 54 (58.1%) were cured after dual therapy of vancomycin plus metronidazole, 7 (7.5%) died of underlying diseases aggravated with CDAD, and 3 (3.2%) were transferred to other hospitals for personal reasons.

Conclusion: The incidence of nosocomial CDAD in China was high. Some risk factors could predispose CDAD.

KEY WORDS: Antibiotic-associated diarrhea, *Clostridium difficile*, Incidence, Risk factors.

doi: <http://dx.doi.org/10.12669/pjms.323.9400>

How to cite this:

Li Y, Huang Y, Li Y, Nie Y. Clinical characteristics of *Clostridium difficile*-associated diarrhea among patients in a tertiary care center in China. Pak J Med Sci. 2016;32(3):736-741. doi: <http://dx.doi.org/10.12669/pjms.323.9400>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Clostridium difficile (*C. difficile*)-associated diarrhea (CDAD) is a common cause of infectious diarrhea in hospitals, which usually occurs as a compli-

cation of antibiotic therapy. CDAD is becoming a growing worldwide health threat with an incidence of approximately 100 per million and a mortality of 1%–2.5% in western countries.¹ Thus, controlling the spread of this infection is urgently needed.

The clinical spectrum of cases presented with CDAD can be extensive, which range from asymptomatic carriage to mild self-limiting diarrhea and more severe pseudo-membranous colitis. Accurate diagnosis of CDAD is crucial in managing individual patients. Over the past two decades, rapid diagnostic methods through testing *C. difficile* toxin (CDT) were developed to detect *C. difficile* infection. Currently, commercial assays to detect both CDT A and B are available with reasonable sensitivity and specificity. Obtaining information of the *C. difficile* infection worldwide is crucial, following the dramatically increasing rate of CDAD and the recent emergence of the new

1. Yongqiang Li,
2. Yi Huang,
3. Yuyuan Li,
4. Yuqiang Nie,

1-4: Department of Gastroenterology, Guangzhou Digestive Diseases Center, Guangzhou First People's Hospital, Guangzhou Medical University, Guangdong, China.

Correspondence:

Prof. Yuqiang Nie,
Department of Gastroenterology,
Guangzhou Digestive Diseases Center,
Guangzhou First People's Hospital, Guangzhou Medical University,
No.#1 Panfu Road, Guangzhou, China. 510180.
Email: nieyq@medmail.com.cn

* Received for Publication: November 20, 2015

* Accepted for Publication: April 12, 2016

highly virulent strains of *C. difficile* in Canada, USA, and Europe.²⁻⁵ However, the data in China are not well documented. In present study, we studied the incidence, risk factors, and clinical characteristics of CDAD in Chinese patients.

METHODS

The study included hospitalized patients admitted to this hospital between April 1, 2008 and March 31, 2010. The patients who exhibited diarrhea after being administered antibiotics for at least 3 days were selected according to the diagnostic criteria issued by Health Ministry of China,⁶ which was adapted from the guideline of American College of Gastroenterology.⁷ These patients suffered from antibiotic-associated diarrhea (AAD) and were included in this study. The diagnosis of AAD was based on the clinical manifestations, i.e., abdominal cramps, profuse diarrhea (bowel movements > three times/day with mucoid, greenish, foul-smelling, and watery stools or pseudo-membrane), low-grade fever, and leukocytosis, which presented several days after initiating antibiotic therapy.

Other gastrointestinal diseases, e.g., bacterial and amebic dysentery, typhoid fever, food poisoning, inflammatory bowel disease, irritable bowel syndrome, lactose intolerance, and colorectal cancer must be ruled out before diagnosis. Stool examinations and colonoscopy were needed when the diagnosis was suspected. Fecal specimens were collected from each patient for the *C. difficile* toxin assay. The patient was diagnosed with CDAD when the result of the assay was positive. A case was considered complicated if one or more of the following was observed: megacolon, perforation, colectomy, shock requiring vasopressor therapy, or death within 30 days after diagnosis. The cases of AAD without CDAD were set as the control. With a ratio of 1:3 for each CDAD cases, 279 matched patients (age \pm 5 years and same gender) from the same department who received antibiotics for at least three days, but had no diarrhea during the same period of time were selected as another control group. The study was approved by the Ethics Committee of the hospital. Written consent was obtained from each participant.

Interview and physical examination: The study was prospectively designed. Face-to-face interview and physical examination were carried out for each subject by specially trained post-graduate students of Guangzhou Medical University and supervised by experienced investigators. Standard questionnaires including demographic data,

current medication use, medical history, and health-relevant behaviors, i.e., alcohol consumption, smoking habits, and dietary habits, were recorded. The patients were followed up for an interval of three days during their stay at the hospital, and their clinical data from patient charts and hospital computer databases were collected for analysis.

Detection of *C. difficile*: Fresh fecal specimens were collected and sent to the laboratory within 2 hours. The tests for *C. difficile* toxin A and B were performed immediately by utilizing enzyme-linked fluorescent assay kits (bioMérieux, France). The tests were carried out according to the instruction of the kits (bioMérieux mini-VIDAS standard). The stool specimens were diluted and centrifuged. The supernatant fluid was placed in the holes of testing kits with CDAD reagent strips inside. The results were measured by bioMérieux equipment. The positive threshold value was set at 0.13. A value < 0.13 was considered negative for CDAD, which suggested AAD. Meanwhile, a value of \geq 0.13 was considered positive, which may imply CDAD.

Variables of observation: The outcomes of interest included the following variables: 1. general conditions: temperature, respiratory rate, heart rate, blood pressure, and consciousness; 2. clinical manifestation: severity and frequency of abdominal pain, frequency of bowel movements, appearance of stools, and abdominal tenderness; 3. laboratory parameters: routine analysis of blood, urine, and stools, liver and kidney function tests, C-reaction protein, and blood gas analysis; 4. administration of medicines: antibiotics, corticosteroid, and immunosuppressive agents; 5. past history and underlying diseases; and 6. therapeutic intervention: nasogastric tube-feeding, urethral catheterization, and tracheal cannula. The Acute Physiology and Chronic Health Evaluation II (APACHE II) score system was employed to further evaluate the severity of the disease.⁸

Statistical analysis: The data were analyzed with SPSS 17.0 (Chicago, IL, USA). The continuous data were expressed as mean \pm SD and examined by Student's t-test. Categorical variables were presented as a percentage and examined by χ^2 tests and Fisher's tests. Statistical significance was set at $P < 0.05$ (two-tailed). Exposure ratio comparison and multivariate regression analyses were performed to evaluate the risk factors.

RESULTS

This general hospital with 1,400 patient beds is located at downtown of Guangzhou, a large city

in southern China with population of 10,045,800. Only patients with advanced diseases are usually admitted to the hospital because of the shortage of medical resources. Between April 1, 2008 and March 31, 2010, 56,172 patients were admitted, of which 39,882 (71.0%) used antibiotics for at least three days. A total of 470 were diagnosed with AAD, among which 93 had CDAD (males, 61; females, 32; aged 68±16 years old) and 377 had AAD without CDAD (males, 211; females, 166; aged 67±19 years old). The incidence of nosocomial CDAD was 166 per million in this hospital. The proportions of AAD and CDAD among inpatients receiving antibiotics were 11.8% and 0.23%, respectively. The proportion of CDAD in AAD was 19.8%. The hospital does not serve a well-defined population. Thus, we were unable to calculate the population-based incidence.

Clinical characteristics of CDAD: Among total 93 patients with CDAD, 70 (75.3%) were elderly (>60 years old). The main underlying diseases of the patients included respiratory failure with 23 cases (24.7%), heart failure with 14 cases (15.1%), cerebrovascular accident with 12 cases (12.9%), gastrointestinal diseases with 16 cases (17.2%),

leukemia with 11 cases (11.8%), post-abdominal operation with 8 cases (8.6%), and others with 9 cases (9.7%). Seventy-one of the patients (76.3%) exhibited concurrent diseases, such as infection, chronic obstructive lung disease, diabetes mellitus, hypertension, heart insufficiency, renal insufficiency, and malnutrition. Forty-two (45.2%) of the patients had more than one concurrent disease. Forty patients (43%) had nasogastric feeding. The average APACHE scores of CDAD patients upon diagnosis were 16.6±4.3. Diarrhea occurred 4–39 days after antibiotic intake. The frequencies of bowel movement were 4–10 per day. Liquid stools were observed in 39 cases (41.9%); mushy stools in 27 cases (29.0%); mucous stools in 16 cases (17.2%), and bloody purulent stools in 11 cases (11.8%), among which pseudo-membranes were found in three cases (3.2%).

Compared with patients with AAD, the patients with CDAD presented significantly longer durations of hospitalization and antibiotic intake, higher serum level of C-reactive protein, lower level of serum albumin, and higher scores of APACHE II and comorbidity (mostly in the lung, heart, liver, and kidney) (Table-I) ($P < 0.05$). The species and numbers of antibiotics were not significantly different between the two groups ($P > 0.05$) except glucopeptide (mainly vancomycin) and nitroimidazole (mainly metronidazole), which protected AAD patients from progressing to CDAD ($P < 0.05$) (Table-I). The multivariate regression logistic analysis demonstrated that C-reactive protein, duration of hospitalization, APACHE II scores, and comorbidity were risk factors (Table-II).

Compared with those of the controls, the variables significantly predisposing CDAD were advanced age, high WBC, nasogastric feeding, high C-reactive protein, high serum albumin level, comorbidity, and APACHE II scores ($P < 0.05$) (Table-III). The multivariate regression logistic analysis showed that nasogastric feeding, C-reactive protein,

Table-I: Clinical characteristics of CDAD patients compared with AAD.

| Variables | CDAD | AAD | P value |
|---------------------------------|---------------|--------------|---------|
| N | 93 | 377 | |
| Age (yrs) | 68 ± 16 | 67 ± 19 | NS* |
| Duration of hospitalization | 18.5 ± 10.4 | 15.2 ± 13.4 | < 0.05 |
| Duration of antibiotics | 10.9 ± 7.1 | 8.8 ± 6.3 | < 0.05 |
| Nasogastric feeding (n) | 40 (45%) | 142 (37.6%) | NS |
| Comorbidity (n) | 73 (78.5%) | 150 (39.8%) | < 0.05 |
| WBC (×10 ¹² /l) | 9.1 ± 3.4 | 8.6 ± 4.2 | NS |
| C-reactive protein (mg/l) | 34.4 ± 35.2 | 23.4 ± 25.8 | < 0.05 |
| Creatinine (umol/l) | 151.9 ± 113.4 | 132.7 ± 25.6 | NS |
| Albumin (g/l) | 29.3 ± 5.4 | 32.0 ± 1.8 | < 0.05 |
| APACHE II scores | 16.6 ± 4.4 | 11.5 ± 5.3 | < 0.05 |
| Intake of antibiotics | | | |
| Third generation cephalosporin | 50 (53.8%) | 175 (46.4%) | NS |
| Second generation cephalosporin | 20 (21.5%) | 80 (21.2%) | NS |
| Quinolones | 39 (41.9%) | 144 (38.2%) | NS |
| Macrolide | 20 (21.5%) | 77 (20.4%) | NS |
| Aminoglycoside | 25 (26.9%) | 88 (23.3%) | NS |
| Carbapenems | 15 (16.1%) | 58 (15.4%) | NS |
| Glucopeptide# | 2(2.2%) | 36 (9.5%) | < 0.05 |
| Nitroimidazole+ | 10(10.8%) | 81 (21.5%) | < 0.05 |
| Quantity of antibiotics | | | |
| Mono | 20 (21.5%) | 86 (22.8%) | NS |
| Dual | 58 (62.4%) | 220 (58.4%) | NS |
| Triple | 15 (16.1%) | 71 (18.8%) | NS |

CDAD: C difficile associated diarrhea;

AAD: Antibiotic-associated diarrhea.

*NS: not significant; # including vancomycin;

+ including metronidazole, ornidazole.

Table-II: Multivariate regression logistic analysis for CDAD in AAD patients.

| Variables | RR | 95% CI | P value |
|---|------|---------------|---------|
| Duration of hospitalization | 2.40 | (1.59, 3.63) | < 0.01 |
| Duration of antibiotics | 0.91 | (0.64, 1.29) | 0.603 |
| C-reactive protein | 2.73 | (1.58, 4.7) | < 0.01 |
| Albumin | 1.34 | (0.81, 2.21) | 0.250 |
| Comorbidity | 5.52 | (3.23, 9.44.) | < 0.01 |
| APACHE II scores | 6.53 | (3.65, 11.68) | < 0.01 |
| Usage of glucopeptide or nitroimidazole | 3.00 | (1.59, 5.79) | < 0.01 |

* RR: relative risk; CI: confidence interval.

Table-III: Clinical characteristics of CDAD patients compared with controls.

| Variables | CDAD | Controls | P value |
|----------------------------|---------------|-------------|---------|
| N | 93 | 279 | |
| Nasogastric feeding (n) | 40 (45%) | 42 (15.1%) | <0.05 |
| Age (yrs) | 68 ± 16 | 59.7 ± 22.6 | <0.05 |
| Comorbidity (n) | 73 (78.5%) | 117 (41.9%) | <0.05 |
| WBC (×10 ¹² /l) | 9.1 ± 3.4 | 8.37 ± 2.8 | <0.05 |
| C-reactive protein (mg/l) | 34.4 ± 35.2 | 13.8 ± 15.5 | <0.05 |
| Creatinine (umol/l) | 151.9 ± 113.4 | 94.1 ± 75.1 | <0.05 |
| Albumin (g/l) | 29.3 ± 5.4 | 34.5 ± 5.7 | <0.05 |
| PACHE II scores | 16.6 ± 4.3 | 9.7 ± 4.4 | <0.05 |

APACHE II, serum albumin level, and comorbidity were risk factors for CDAD (Table-IV).

The impact of antibiotics selection on CDAD: Compared with those of the controls, the administration of the third generation cephalosporin, quinolones significantly increased the incidence of CDAD after administering the third generation cephalosporin. However, glucopeptide and nitroimidazole significantly decreased the incidence of CDAD (Table-V).

Prognosis of patients with CDAD: Discontinuation of antibiotics was possible in 49 patients. Ninety-three patients with CDAD were given 0.25–0.5 g Vancomycin four times a day. After the 7 day treatment, 29 patients (31.2%) were cured. The poor responders were administered 200–400 mg Metronidazole four times a day. The remaining 54 patients (58.1%) were cured with dual therapy, among which 38 (40.9%) responded within the next 7 days and 11 responded within 27 days. Seven patients (7.5%) died of underlying diseases aggravated with CDAD. Three patients (3.2%) were referred to other hospitals for personal reasons.

DISCUSSION

There are several laboratory techniques to identify *C. difficile* infection. However, the best standard method has not been clearly established. *C. difficile* cytotoxin neutralization and toxigenic culture are

Table-V: Antibiotics intake in CDAD and control groups.

| Antibiotics | CDAD | Control | χ^2 | P |
|-------------------------------------|------------|-------------|----------|-------|
| Used antibiotics | | | | |
| Third generation cephalosporin (n) | 50 (53.8%) | 96 (34.4%) | 10.96 | <0.05 |
| Second generation cephalosporin (n) | 20 (21.5%) | 76 (27.2%) | 1.20 | NS |
| Quinolones (n) | 39 (41.9%) | 74 (26.5%) | 7.83 | <0.05 |
| Macrolide (n) | 20 (21.5%) | 80 (28.7%) | 1.82 | NS |
| Aminoglycoside (n) | 25 (26.9%) | 60 (21.5%) | 1.14 | NS |
| Carbopenems | 15 (16.1%) | 24 (8.6%) | 4.21 | <0.05 |
| Glucopeptide (n) | 2(2.2%) | 27 (9.7%) | 5.50 | <0.05 |
| Nitroimidazole (n) | 10(10.8%) | 74 (26.5%) | 9.92 | <0.05 |
| Number of antibiotics | | | | |
| Mono | 20 (21.5%) | 75 (26.8%) | 1.06 | NS |
| Dual | 58 (62.4%) | 176 (63.1%) | 0.02 | NS |
| Triple | 15 (16.1%) | 28 (10.0%) | 2.53 | NS |

* NS: >0.05

usually considered as the primary reference tests. However, these tests are time consuming and require equipment and expertise. Stool culture alone is not recommended because not all *C. difficile* strains are toxigenic. According to the new guidelines published by AJG in 2013,⁹ Nucleic acid amplification tests (NAAT) such as polymerase chain reaction (PCR) for *C. difficile* toxin genes appear to be sensitive and specific and may be used as a standard diagnostic test for CDI. *C. difficile* glutamate dehydrogenase (GDH) test is a sensitive test with low specificity. This test is applied as a screen test to eliminate most samples (that test negative) for further testing.¹⁰ GDH may be used in association with toxin A and B enzyme immunoassay (EIA) testing. EIA for toxin A/B used to be the most widely used diagnostic test because it is fast, inexpensive, and has high specificity. Although its sensitivity is lower than that of NAAT, EIA can identify the toxigenic strains, which are of clinical importance.¹¹ In this study, we focused on the pathogenic strains of *C. difficile*. Thus, EIA for *C. difficile* toxin A/B was chosen. Moreover, this testing approach was suggested by UK consensus at that time.¹²

Table-IV: Multivariate regression logistic analysis for the risk of CDAD in inpatients.

| Variables | β | SE | Wald | RR (95% CI) | P value |
|--------------------------------|---------|-------|--------|--------------------|---------|
| Nasogastric feeding | 1.449 | 0.268 | 29.20 | 4.26 (2.52,7.20) | < 0.05 |
| Comorbidity | 1.620 | 0.281 | 33.47 | 5.05 (2.92,8.75) | < 0.05 |
| APACHE II | 2.44 | 0.408 | 35.91 | 11.52 (5.18,25.63) | < 0.05 |
| C-reactive protein | 0.938 | 0.257 | 13.36 | 2.55 (1.55,4.22) | < 0.05 |
| Albumin | 0.801 | 0.247 | 10.52 | 2.23 (1.37,3.6) | < 0.05 |
| Numbers of antibiotics | 0.294 | 0.286 | 1.056 | 1.34 (0.766,2.352) | 0.304 |
| Third generation cephalosporin | 0.779 | 0.243 | 10.257 | 2.18(1.35,3.51) | < 0.05 |
| Quinolones | 0.679 | 0.250 | 7.357 | 1.97(1.21,3.22) | < 0.05 |
| Glucopeptide | -1.596 | 0.743 | 4.615 | 0.203(0.05,0.87) | < 0.05 |
| Nitroimidazole | -1.112 | 0.361 | 9.475 | 0.33(0.16,0.67) | < 0.05 |

CDAD became prevalent in the 1960s and 1970s with the introduction of broad-spectrum antibiotics to the clinical practice. The incidence of *C. difficile* infection increased rapidly. During the middle and late 1990s, the reported incidence of *C. difficile* infection in acute care hospitals in the United States remained stable at 30 to 40 cases per million in population. In 2001, this rate increased to almost 50, with subsequent increases to 92.8 in 2011². The outbreaks of *C. difficile* infection in 2003 with increased severity of illness were reported in Quebec, Canada with an incidence of 156.3 per million, and 102.0 to 866.5 among patients aged 65 years old or more. The outbreaks were caused by a new virulent strain (called PCR ribotype 027).¹³ Later, many parts of the United States reported similar outbreaks caused by the same strain;^{14,15} the outbreaks spread to Europe afterward.¹⁶ Reported mortality rates from CDAD in the United States increased from 5.7 per million in 1999 to 23.7 per million in 2004. The rate increased to 1%–2.5% in 2008 because of the emergence of a highly virulent strain of *C. difficile*.¹ The incidence and mortality rate of *C. difficile* infection in other places, particularly in developing countries, have not been documented. The present study reveals a high incidence of nosocomial CDAD (166 per million) at the hospital in Guangzhou, China. Although we were unable to calculate population-based incidence and determine the strain of *C. difficile*, these results should warrant more attention in the investigation of CDAD.

Identifying patients who are at high risk for CDAD early in the course of their infection may help improve outcomes, although the predictors are not well known. Compared with AAD patients, CDAD patients usually present more severe clinical manifestations and have more concurrent illnesses.¹⁷ Laboratory markers, such as leukocytosis, increased creatinine levels, and decreased albumin levels, were reported to correlate with the development and poor outcome for patients with CDAD.¹⁸ Disrupting normal intestinal flora by antibiotics is a well-known mechanism for CDAD. Thus, antibiotic intake is the most important risk factor. Other drugs, such as immunosuppressive agents and proton pump inhibitors, are also significant risk factors for CDAD precipitation. Although the antibiotics most frequently implicated in the predisposition to *C. difficile* infection are fluoroquinolones, clindamycin, cephalosporins, and penicillins, virtually all antibiotics (including metronidazole and vancomycin) can predispose the patient to *C. difficile* infection. Risk is higher when the patients are on

multiple antibiotics and undergo longer courses of therapy.¹⁹ In this study, no difference was observed in most antibiotic intakes between AAD and CDAD groups. The administration of glucopeptide (mainly vancomycin) and nitroimidazole were even negatively related to CDAD (as protective factors). In non-drug risk factors, nasogastric feeding, comorbidity, elevated serum C-reactive protein, and creatinine levels, decreased serum albumin, and increased APACHE II scores were all risk factors of developing CDAD in comparison with that of the controls. However, compared with that of AAD, only comorbidity, elevated serum C-reactive protein, duration of hospitalization, and increased APACHE II scores were the risk factors of CDAD. This group of patients shared most non-drug risk factors of those reported in developed countries. This outcome implied that the host responses to the infection of Chinese and their western counterparts were identical.

The usual treatment for CDAD was to stop administration of antibiotics meant for other purposes and to immediately start treatment with metronidazole or vancomycin. Patients who remain on antibiotics while undergoing CDAD treatment are likely to experience metronidazole treatment failure.²⁰ In this study, administering glucopeptide (mainly vancomycin) and nitroimidazole was a protective factor, and most patients were successfully cured with these two kinds of drugs probably because patients were infected with the strain of *C. difficile* sensitive to the drugs. The resistant strain of *C. difficile* seemed uncommon in China.

In conclusion, to our knowledge, the present work is a rare prospective study that investigates the incidence and clinical characteristics of CDAD in mainland of China.^{21–25} We determined a high incidence of *C. difficile* infection in China. The major limitations of this study were the comparatively small sample size and the testing method. This work still provides vital information, which may provide basis for future research focusing on the prevention of CDAD.

Declaration of interest: The authors declare that they have no conflict of interest.

REFERENCES

1. Zilberberg MD, Shorr AF, Kolfel MH. Increase in adult *Clostridium difficile*-related hospitalizations and case-fatality rate, United States, 2000–2005. *Emerg Infect Dis.* 2008;14:929–931. doi:10.3201/eid1406.071447

2. Lessa FC, Mu Y, Bamberg WM, Beldavs ZG, Dumyati GK, Dunn JR, et al. Burden of Clostridium difficile infection in the United States. *N Engl J Med.* 2015;372:825-834. doi:10.1056/NEJMoa1408913
3. Kelly CP, LaMont JT. Clostridium difficile--more difficult than ever. *N Engl J Med.* 2008;359:1932-1940. doi:10.1056/NEJMra0707500
4. Kuijper EJ, Coignard B, Tull P, difficile ESGfC, States EUM, European Centre for Disease P, et al. Emergence of Clostridium difficile-associated disease in North America and Europe. *Clin Microbiol Infect.* 2006;12(Suppl 6):2-18. doi:10.1111/j.1469-0691.2006.01580.x
5. Muto CA, Pokrywka M, Shutt K, Mendelsohn AB, Nouri K, Posey K, et al. A large outbreak of Clostridium difficile-associated disease with an unexpected proportion of deaths and colectomies at a teaching hospital following increased fluoroquinolone use. *Infect Control Hosp Epidemiol.* 2005;26:273-280. doi:10.1086/502539
6. China HMo. Diagnostic criteria for nosocomial infection. Health Ministry of China, Beijing 2001:10-12.
7. Fekety R. Guidelines for the diagnosis and management of Clostridium difficile-associated diarrhea and colitis. American College of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol.* 1997;92(5):739-750.
8. LeGall JR, Loirat P, Alperovitch A. APACHE II--a severity of disease classification system. *Crit Care Med.* 1986;14:754-755.
9. Surawicz CM, Brandt LJ, Binion DG, Ananthakrishnan AN, Curry SR, Gilligan PH, et al. Guidelines for diagnosis, treatment, and prevention of Clostridium difficile infections. *Am J Gastroenterol.* 2013;108:478-498; quiz 499. doi:10.1038/ajg.2013.4
10. DuPont HL, Garey K, Caeiro JP, Jiang ZD. New advances in Clostridium difficile infection: changing epidemiology, diagnosis, treatment and control. *Curr Opin Infect Dis.* 2008;21:500-507. doi:10.1097/QCO.0b013e32830f9397
11. Debast SB, van Kregten E, Oskam KM, van den Berg T, Van den Berg RJ, Kuijper EJ. Effect on diagnostic yield of repeated stool testing during outbreaks of Clostridium difficile-associated disease. *Clin Microbiol Infect.* 2008;14:622-624. doi:10.1111/j.1469-0691.2008.01999.x
12. Vonberg RP, Kuijper EJ, Wilcox MH, Barbut F, Tull P, Gastmeier P, et al. Infection control measures to limit the spread of Clostridium difficile. *Clin Microbiol Infect.* 2008;14(Suppl 5):2-20. doi:10.1111/j.1469-0691.2008.01992.x
13. Pepin J, Valiquette L, Alary ME, Villemure P, Pelletier A, Forget K, et al. Clostridium difficile-associated diarrhea in a region of Quebec from 1991 to 2003: a changing pattern of disease severity. *CMAJ.* 2004;171:466-472. doi:10.1503/cmaj.1041104
14. McDonald LC, Killgore GE, Thompson A, Owens RC, Jr., Kazakova SV, Sambol SP, et al. An epidemic, toxin gene-variant strain of Clostridium difficile. *N Engl J Med.* 2005;353:2433-41 doi:10.1056/NEJMoa051590
15. Loo VG, Poirier L, Miller MA, Oughton M, Libman MD, Michaud S, et al. A predominantly clonal multi-institutional outbreak of Clostridium difficile-associated diarrhea with high morbidity and mortality. *N Engl J Med.* 2005;353:2442-2449. doi:10.1056/NEJMoa051639
16. Kuijper EJ, Coignard B, Brazier JS, Suetens C, Drudy D, Wiuff C, et al. Update of Clostridium difficile-associated disease due to PCR ribotype 027 in Europe. *Euro Surveill.* 2007;12:E1-2.
17. Barbut F, Gariazzo B, Bonne L, Lalande V, Burghoffer B, Luiuz R, et al. Clinical features of Clostridium difficile-associated infections and molecular characterization of strains: results of a retrospective study, 2000-2004. *Infect Control Hosp Epidemiol.* 2007;28:131-139. doi:10.1086/511794
18. Moshkowitz M, Ben-Baruch E, Kline Z, Shimoni Z, Niven M, Konikoff F. Risk factors for severity and relapse of pseudomembranous colitis in an elderly population. *Colorectal Dis.* 2007;9:173-177. doi:10.1111/j.1463-1318.2006.01013.x
19. Drudy D, Harnedy N, Fanning S, Hannan M, Kyne L. Emergence and control of fluoroquinolone-resistant, toxin A-negative, toxin B-positive Clostridium difficile. *Infect Control Hosp Epidemiol.* 2007;28:932-940. doi:10.1086/519181
20. Modena S, Gollamudi S, Friedenber F. Continuation of antibiotics is associated with failure of metronidazole for Clostridium difficile-associated diarrhea. *J Clin Gastroenterol.* 2006;40:49-54.
21. Wang X, Cai L, Yu R, Huang W, Zong Z. ICU-Onset Clostridium difficile infection in a university hospital in China: a prospective cohort study. *PLoS One.* 2014;9:e111735. doi:10.1371/journal.pone.0111735
22. Huang H, Wu S, Chen R, Xu S, Fang H, Weintraub A, et al. Risk factors of Clostridium difficile infections among patients in a university hospital in Shanghai, China. *Anaerobe.* 2014;30:65-69. doi:10.1016/j.anaerobe.2014.08.015
23. Lv Z, Peng GL, Su JR. Factors associated with Clostridium difficile diarrhea in a hospital in Beijing, China. *Braz J Med Biol Res.* 2014;47:1085-90
24. Jin K, Wang S, Huang Z, Lu S. Clostridium difficile infections in China. *J Biomed Res.* 2010;24:411-416. doi:10.1016/S1674-8301(10)60055-3
25. Huang H, Wu S, Wang M, Zhang Y, Fang H, Palmgren AC, et al. Molecular and clinical characteristics of Clostridium difficile infection in a University Hospital in Shanghai, China. *Clin Infect Dis.* 2008;47:1606-1608. doi:10.1086/593365

Authors' Contributions:

Yuqiang Nie: Conceived and designed the experiments.

Yongqiang Li, Yi Huang: Performed the experiments.

Yongqiang Li, Yi Huang, Yuyuan Li: Analyzed the data.

Yongqiang Li, Yi Huang, Yuqiang Nie: Contributed reagents/materials/analysis tools.

Yongqiang Li, Yuqiang Nie: Prepared the manuscript.